

## Research paper

## Testing post-IR IRSL luminescence dating methods in the southwest Mojave Desert, California, USA

Andrew S. Carr<sup>a,\*</sup>, Alex S. Hay<sup>a,\*\*</sup>, D. Mark Powell<sup>a</sup>, Ian Livingstone<sup>b</sup><sup>a</sup> School of Geography, Geology and the Environment, University of Leicester, University Road, Leicester LE1 7RH, UK<sup>b</sup> The Graduate School, University of Northampton, Northampton, NN2 7AL, UK

## A B S T R A C T

The Mojave Desert presents an array of Pleistocene lacustrine deposits and aeolian landforms to which, at times, it has proved challenging to apply luminescence methods. We tested the suitability of K-feldspar post-IR IRSL methods using two sites with independent radiocarbon dating – shorelines at Harper Lake and Silver Lake – considering: 1) overall performance of the post-IR IRSL 225 °C (pIRIR<sub>225</sub>) protocol; 2) effect of test dose size on pIRIR<sub>225</sub> D<sub>e</sub>; 3) anomalous fading correction of pIRIR<sub>225</sub> ages; 4) preliminary single grain pIRIR<sub>225</sub> results.

We observe consistently good performance of the single aliquot pIRIR<sub>225</sub> protocol, with good dose recovery, acceptable recycling ratios, low recuperation and low inter-aliquot scatter. The pIRIR<sub>225</sub> ages for Silver Lake (8.8 ± 0.4 and 11.3 ± 0.5 ka) and Harper Lake (both 25.4 ± 1.4 ka) are in substantially better agreement with the independent dating than low temperature (50 °C) IRSL and quartz OSL ages. pIRIR<sub>225</sub> fading rates are reduced to ~2.0–2.5% per decade, but there remains a tendency for under-estimation when using uncorrected ages. A need for fading correction is further implied at Harper Lake via comparison with multi-elevated temperature (MET)-PIR age plateaus and pIRIR<sub>290</sub> measurements, although at the younger Silver Lake site these methods produce ages nearly identical to the uncorrected pIRIR<sub>225</sub> ages. Preliminary single grain pIRIR<sub>225</sub> measurements suggest a ~25–30% usable grain yield. At Silver Lake the single grain and single aliquot ages agree well despite over-dispersion of the single grain equivalent dose distribution. At Harper Lake the single grain and single aliquot pIRIR<sub>225</sub> ages also agree well, although a population of insensitive, lower D<sub>e</sub> grains is observed. These grains are not associated with significantly higher fading rates.

## 1. Introduction

The Mojave Desert (southwest USA) preserves abundant evidence for Pleistocene palaeo-lakes (Enzel et al., 2003) and relict aeolian deposits (Lancaster and Tchakerian, 1996). Various luminescence dating methods have been applied, including quartz optically stimulated luminescence (OSL) (Bateman et al., 2012; Fuchs et al., 2015), K-feldspar thermoluminescence (TL) and infra-red stimulated luminescence (IRSL; Clarke et al., 1995; Rendell and Sheffer, 1996). Presently there are conflicting ages between studies (e.g. Rendell and Sheffer, 1996; Bateman et al., 2012), within sites (stratigraphic inversions), and contrasting results compared to independent dating (e.g. Owen et al., 2007). Quartz may be an unreliable dosimeter in the Mojave due to its low sensitivity and a contaminating K-feldspar signal (Lawson et al., 2012). K-feldspar is, however abundant in Mojave sediments and is a potentially advantageous mineral choice given the relatively high environmental dose rates (typically > 3 Gy ka<sup>-1</sup>). Previous applications

of K-feldspar IRSL (Rendell and Sheffer, 1996) did not include anomalous fading correction and subsequent studies using low temperature IRSL have reported variable, but sometimes high fading rates (Garcia et al., 2014).

We consider the reliability of K-feldspar ages derived via post-IR IRSL (pIRIR) methods, which can isolate a slower (or non) fading IRSL signal (Thomsen et al., 2008; Buylaert et al., 2012). Demonstrating the suitability of pIRIR approaches would be an important step in improving chronological control in the Mojave, and recent applications have shown promise (McGuire and Rhodes, 2015; Roder et al., 2012). We sampled two sites with independent dating to consider pIRIR protocol performance.


## 2. Study sites

Two palaeo-lakes in the Mojave River catchment were analysed; Harper Lake and Silver Lake (Fig. 1). Sourced in the San Bernardino

\* Corresponding author.

\*\* Corresponding author.

E-mail addresses: [asc18@le.ac.uk](mailto:asc18@le.ac.uk) (A.S. Carr), [ash24@le.ac.uk](mailto:ash24@le.ac.uk) (A.S. Hay).



**Fig. 1.** A) Map of the Mojave River catchment, showing the major palaeo-lakes and the present Mojave River course. B) Site stratigraphy with previously published (re-calibrated) radiocarbon dates and luminescence ages for Harper Lake. C) Silver Lake stratigraphy, with quartz OSL ages and re-calibrated radiocarbon dates (Table S2) for LFA6 and LFA8 (modified after Owen et al., 2007).

Mountains to the southwest, the Mojave River experienced episodes of perennial flow during the Pleistocene, periodically maintaining Lake Manix, Harper Lake and the downstream Silver Lake. This catchment history is discussed elsewhere (Meek, 1999; Enzel et al., 2003; Wells et al., 2003; Reheis et al., 2012, 2015).

Silver Lake formed part of pluvial Lake Mojave (Wells et al., 2003) (Fig. 1a; 1c). Several sites demonstrate lake high-stands (following Wells et al., 2003) at ~22–19 cal ka BP (18.4–16.6 ka; “Lake Mojave 1”) and ~16.6–13.3 cal ka BP (13.7–11.4 ka; “Lake Mojave 2”), with intermittent inundation at 13.3–9.8 cal ka BP (~11.4–8.7 ka). A spit-shoreline complex at “Silver Quarry” (Ore and Warren, 1978) was subject to a detailed investigation combining radiocarbon dating of the freshwater bivalve *Anodonta californiensis*, quartz OSL and fine-grain K-feldspar IRSL Multi-Aliquot Additive Dose (MAAD) methods (Owen et al., 2007). Our luminescence samples were obtained from Owen et al.’s (2007) Lithofacies Associations (LFA) LFA8 (SL14-1; 0.4 m) and LFA6 (SL14-2; 1.2 m) (Fig. 1c). Using the BCal Bayesian analysis software (Table S2), Owen et al. (2007) assigned age ranges of 12.1–11.6 cal ka BP to LFA 8 (7 dates) and 12.2–12.5 cal ka BP to LFA 6 (2 dates). Their quartz OSL ages for LFA 8 (SL125) and LFA6 (SL126) were  $6.6 \pm 0.7$  ka and  $6.5 \pm 0.6$  ka respectively.

Harper Basin (Fig. 1a) is presently isolated from the Mojave River, but was likely fed by periodic Mojave River avulsions (Meek, 1999). Lake beds are exposed at ‘Mountain View Hill’ where radiocarbon dates

from *A. californiensis* shells of 24,055–33,059 cal yr. BP ( $24,440 \pm 2190$   $^{14}\text{C}$  yr BP) and 28,375–29,790 cal yr. BP ( $25,000 \pm 310$   $^{14}\text{C}$  yr BP) were first reported (with a third infinite age; Meek, 1999). Garcia et al. (2014) presented eight new *A. californiensis* radiocarbon dates and luminescence ages from coarse-grain (125–150  $\mu\text{m}$ ) post-IR quartz SAR and fine grain (4–11  $\mu\text{m}$ ) K-feldspar MAAD IRSL (Fig. 1b). The new radiocarbon dates ranged from 33,410 to 39,788, cal yr. BP; Table S2; Fig. 1b), with fading-corrected IRSL ages of  $28 \pm 2$  ka to  $46 \pm 3$  ka (7.2% per decade fading rate). The quartz ages were substantially younger (17–19 ka). Garcia et al. (2014) argued for a probable age of 40–45 ka, but there is variability within and between the radiocarbon and IRSL ages, with the former close to the limits of the method. The independent dating control at Harper Lake is thus less firm than Silver Lake. We sampled the same section and took samples from the beach unit (Fig. 5 of Garcia et al., 2014) at 0.84 and 1.25 m (Fig. 1c).

### 3. Methods

180–250  $\mu\text{m}$  quartz and K-feldspar grains were isolated, with K-rich feldspars obtained via density separation at  $2.58 \text{ g cm}^{-3}$  and etched in 10% HF for 10 min. All samples were analysed on a Risoe DA20 TL/OSL reader, with quartz luminescence detected through a Hoya U340 filter and IRSL through Schott BG39 and Corning 7–59 filters. Quartz

**Table 1**  
Equivalent doses and associated age estimates for the various K-feldspar IRSL methods and the quartz SAR.

Sample	CAM D <sub>e</sub> IR 50 (Gy)	Age IR50 (ka)	Age IR50 (ka) Fading correct	CAM D <sub>e</sub> pIRIR <sub>225</sub> (Gy) (n)	Age pIRIR <sub>225</sub> (ka) <sup>†</sup>	Age pIRIR <sub>225</sub> (ka) correct	Single grain pIRIR <sub>225</sub> (Gy) (n)	Single grain age-pIRIR <sub>225</sub> (ka)	CAM D <sub>e</sub> pIRIR <sub>290</sub> (Gy) (n)	Age pIRIR <sub>290</sub> (ka) <sup>†</sup>	CAM D <sub>e</sub> Quartz* (Gy) (n)	Age quartz (ka)
SL14-1	25.2 ± 0.95 (20)	6.36 ± 0.30	10.3 ± 0.6	34.9 ± 1.2 (20)	8.8 ± 0.4	10.1 ± 0.5	35.3 ± 3.22 (21)	8.9 ± 0.9	33.1 ± 1.29 (4) <sup>†</sup>	8.4 ± 0.4	16.3 ± 2.0	5.2 ± 0.6
SL14-2	37.4 ± 1.23 (19)	8.26 ± 0.40	12.4 ± 0.8	51.2 ± 1.6 (19)	11.3 ± 0.5	11.9 ± 0.6	nd	nd	nd	nd	nd	nd
HL14-1	54.4 ± 1.91 (22)	13.0 ± 0.70	42.1 ± 20	107 ± 3.5 (22)	25.4 ± 1.4	29.1 ± 1.9	94.0 ± 6.2 (28)	22.4 ± 1.8 <sup>‡</sup>	140 ± 4.8 (8)	33.4 ± 1.9	85.6 ± 13.8 (3)	25.6 ± 4.3
HL14-2	56.9 ± 2.12 (20)	13.9 ± 0.80	34.6 ± 7.8	104 ± 3.2 (20)	25.4 ± 1.4	29.9 ± 2.0	nd	nd	152 ± 7.0 (8)	37.3 ± 2.3	nd	nd

<sup>†</sup>No residual subtraction was performed.

<sup>\*</sup>For Harper Lake the quartz D<sub>e</sub> is derived from three acceptable aliquots using early background subtraction.

<sup>‡</sup># 112.4 ± 6.1 Gy and 26.8 ± 1.9 ka using the brightest 50% of grains.

<sup>‡</sup>Excluding high outlier aliquot with D<sub>e</sub>/age of 42 Gy/10.8 ka (Fig. 3).

equivalent doses on 2 mm aliquots were determined using the single aliquot regeneration (SAR) protocol (Murray and Wintle, 2000) employing post-IR (50 °C) blue LED (125 °C) stimulation. 2 mm aliquots of K-feldspar were analysed using a pIRIR protocol comprising a 50 °C IRSL stimulation and a subsequent 225 °C stimulation (henceforth pIRIR<sub>225</sub>) with a 250 °C preheat (1 min). Anomalous fading rates were determined following Auclair et al. (2003) with corrections following Huntley and Lamothe (2001), using the R package “Luminescence” (Kreutzer et al., 2012).

Dose rates were determined using *in-situ* gamma spectrometry and ICP-MS (Table S1). We used the estimated water contents of Owen et al. (2007) and Garcia et al. (2014) (10 ± 5% and 14.5 ± 5% respectively). A 5% absolute change in water content produces an age difference of 200–300 years at Silver Lake and ~380 years at Harper Lake.

## 4. Results

### 4.1. Silver Lake

pIRIR<sub>225</sub> ages (Table 1 and Fig. S1) were obtained using a moderate (23% of D<sub>e</sub>) test dose. Sample SL14-1 (LFA8) produced a fading-uncorrected pIRIR<sub>225</sub> age of 8.8 ± 0.4 ka, and sample SL14-2 (LFA6) an age of 11.3 ± 0.5 ka. The pIRIR<sub>225</sub> residual D<sub>e</sub>s following 48 h of (UK) daylight were 0.8 and 1.0 Gy. A quartz OSL age of 5.2 ± 0.5 ka for SL14-1 (LFA8) is comparable to that of Owen et al. (2007) and is much younger than the radiocarbon dates. All quartz aliquots are rejected if the fast ratio criterion (average ratio 2.4 ± 1.7) is applied (Durcan and Duller, 2011) and in light of the signal contamination test results (Fig. S2, after Lawson et al., 2012) this age is considered unreliable. We infer this probably also applies to Owen et al.'s (2007) quartz ages.


The pIRIR<sub>225</sub> ages are in better agreement with the radiocarbon dating (BCal ages 12.1–11.6 cal yr. BP and 12.5–12.2 cal yr BP for LFA8 and LFA6). They show low D<sub>e</sub> over-dispersion (3–6%; Fig. S1), good dose recovery (ratios 1.00 ± 0.01 (SL14-1) and 0.99 ± 0.01 (SL14-2)), low recuperation (< 2% for all aliquots) and recycling ratios consistent with unity (e.g. SL14-1 average 1.02 ± 0.02). The SL14-1 fading-uncorrected pIRIR<sub>225</sub> age is younger than the LFA8 BCal radiocarbon age range, although it is within uncertainties of the youngest radiocarbon date from this LFA (AP9; 9081–9322 cal yr BP; Table S2; Fig. S4). SL14-2 is within 2 sigma uncertainties of the LFA6 BCal age (12.2–12.5 ka). Thus, although both pIRIR<sub>225</sub> ages show better agreement with the independent dating, it is prudent to consider possible underestimation relative to the radiocarbon dating.

### 4.2. Test dose size

Test dose size has been shown to impact sample D<sub>e</sub> within pIRIR protocols (Li et al., 2014; Lui et al., 2016; Yi et al., 2016; Colarossi et al., 2018). At Silver Lake the natural D<sub>e</sub> was determined for test doses between 4 and 65% of the expected D<sub>e</sub> (Fig. 2). The results suggest possible age underestimation at low test doses (but note the uncertainties), with a much clearer tendency at high doses. Moderate (23–30% of D<sub>e</sub>) test doses produced ages closest the expected age. Considering the dose response curves (DRC) for low (3.8% of D<sub>e</sub>), moderate (27% of D<sub>e</sub>) and high (65% of D<sub>e</sub>) test doses (Fig. S3), low test dose DRCs saturate faster (D<sub>0</sub> = ~43 Gy compared to > 150 Gy for 27% test dose), with little difference between moderate and high test doses. The age difference between moderate and high test doses seems to reflect a lower Ln/Tn for the latter. The test dose used for the age estimates in Table 1 (23%) is thus unlikely to be a source of age-underestimation.

### 4.3. Ages and fading rates

The pIRIR<sub>225</sub> fading rates are 2.1 ± 0.3% and 0.7 ± 0.3% per decade for SL14-1 and SL14-2 and fading correction brings them into



**Fig. 2.** Relationship between test dose size and sample age for (A) Silver Lake SL14-1 and (B) Harper Lake HL14-1. In A the horizontal line shows the lower 11.6 ka BCal estimate for LFA8 (see Table S2). In B the lowest calibrated (median value) radiocarbon age for HL14-1 is shown with a solid line; (Meek, 1999), with the natural luminescence signal as filled circles. The HL14-1 dose recovery results (open squares) are shown on the secondary Y axis and the dashed line marks a dose recovery ratio of 1. Averages and standard deviations of 3 aliquots are shown.


better (SL14-1) and very good (SL14-2) agreement with the radiocarbon ages (Table 1; Fig. S4). The fading rates for the IR 50 °C are  $6.5 \pm 0.3\%$  and  $5.6 \pm 0.4\%$  for SL14-1 and SL14-2, but with fading-correction ( $10.3 \pm 0.6$  and  $12.4 \pm 0.8$  ka) they show good correspondence with the fading-corrected pIRIR<sub>225</sub> and radiocarbon ages.

The need for fading correction of the post-IR IRSR signal may be removed by using a higher temperature second IR stimulation (pIRIR<sub>290</sub>; Buylaert et al., 2012). This is usually at the expense of a larger unbleached/residual IRSR signal (Kars et al., 2014) and in water-lain deposits, it may be advantageous to utilise a more easily bleached signal (i.e. pIRIR<sub>225</sub>). To assess this further, we compared the pIRIR<sub>225</sub> ages and independent ages with the pIRIR<sub>290</sub> (Buylaert et al., 2012) and multiple elevated temperature (MET) PIR (Li and Li, 2011) methods. For sample SL14-1, we observe possible MET-PIR plateau above 200 °C, but the 250 °C age ( $8.8 \pm 0.5$  ka) matches the fading-uncorrected pIRIR<sub>225</sub> age (Fig. 3). Although pIRIR<sub>290</sub> and MET-300 °C data are broadly within this range, they show more inter-aliquot scatter, perhaps indicating the unsuitability of higher preheating/stimulation temperatures. Increasing the first stimulation temperature (for pIRIR<sub>225</sub>) to 80 °C or 110 °C increases the age of SL14-1 to  $9.9 \pm 0.4$  and  $9.7 \pm 0.4$  ka, perhaps implying removal of a fading-prone signal. However, the

trend does not continue with higher (180 °C) first stimulation temperatures ( $8.7 \pm 0.5$  ka). Thus, the MET-PIR 250 °C age, pIRIR<sub>290</sub> and the uncorrected pIRIR<sub>225</sub> age for SL14-1 all fall at the lower edge of the expected BCal age range (Fig. 3 and S4) and it is presently unclear whether the MET plateau represents a non-faded age.


#### 4.4. Single grain analysis

The single aliquot data show limited inter-aliquot scatter, but given potential signal averaging for K-feldspars (Trauerstein et al., 2014) and the lacustrine context, preliminary single grain measurements were conducted for SL14-1. Grains were mounted on single aliquot disks and stimulated with the IR LED. A dose recovery experiment was conducted, comprising room temperature IR bleaching for 200 s and a 33 Gy dose. Of 96 analysed, 22 grains produced acceptable signals (test dose > 3 sigma above background, recycling ratios between 0.8 and 1.2, recuperation < 5%). The central age model (CAM) dose recovery ratio was  $1.02 \pm 0.03$  (identical to arithmetic mean). The  $D_e$  over-dispersion from the dose recovery experiment was low at  $3.3 \pm 0.4\%$  (cf. Rhodes, 2015; Brown et al., 2015). This OD was added to the individual grain  $D_e$  uncertainties for analysis of the natural  $D_e$ . A natural



**Fig. 3.** MET-PIR plateaus and pIRIR<sub>225</sub> and pIRIR<sub>290</sub> age estimates for samples SL14-1 and HL14-2. The BCal age range is shown for SL14-1 (with the median calibrated age for the youngest sample (AP9) also shown). The range of calibrated radiocarbon ages (median calibrated values in Table S2) are plotted for Harper Lake. Note that the Harper Lake quartz age is from sample HL14-1 (Table 1).





**Fig. 4.** Single grain radial plots (Vermeesch, 2009) for samples SL14-1 and HL14-1. The samples are grey-scaled by sensitivity (first (background subtracted) test dose response in counts  $s^{-1} Gy^{-1}$ ). For Harper Lake the two sigma lines are centred on the CAM estimate for all accepted grains, with the marker line indicating the single aliquot CAM  $D_e$ . Similarly, the marker line for SL1-1 is the single aliquot CAM  $D_e$ .

equivalent dose was derived from 21 grains (of 96 measured). The data show significant ( $37 \pm 6\%$ ) over-dispersion, but the CAM-derived age ( $8.9 \pm 0.9$  ka) is identical to the single aliquot result (Fig. 4). The distribution of grain brightness (Fig. 4 and S5) is skewed (50% of light sum from 18% of grains), but there is no relationship between grain sensitivity and equivalent dose (c.f. Rhodes, 2015), nor is there a correlation between grain fading rates and equivalent dose.

#### 4.5. Harper Lake

Harper Lake produced two identical pIRIR<sub>225</sub> ages (test dose 10% of  $D_e$ ) of  $25.4 \pm 1.4$  ka (Table 1; Fig. S1). The fading-uncorrected ages are within uncertainties of one of Meek's (1999) radiocarbon ages, lower than all other radiocarbon ages, and within uncertainties of one fine-grain IRSL age ( $28 \pm 2$  ka) (Garcia et al., 2014; Fig. S6). pIRIR<sub>225</sub> data show good dose recovery ( $0.98 \pm 0.01$  (HL14-1) and  $0.97 \pm 0.01$  (HL14-2)), good recycling ratios (averages  $1.01 \pm 0.02$  and  $1.02 \pm 0.02$ ) and low recuperation (all aliquots  $< 0.5\%$ ). Residuals following 48 h of daylight were 5.5 and 3.5 Gy. The 50 °C IR ages are  $13.0 \pm 0.7$  ka and  $13.9 \pm 0.8$  ka. Quartz performance was poor (Figs. S2 and S6) with most aliquots rejected for excessive recuperation (average  $\sim 13\%$ ) using late background subtraction. A quartz age for HL14-1 from 3 acceptable aliquots using early background subtraction was  $25.7 \pm 4.4$  ka, and  $23.6 \pm 2.8$  ka for the single acceptable aliquot using late background subtraction.

#### 4.6. Test dose size

For Harper Lake, the effect of test dose size was investigated with a dose recovery experiment and using the natural IRSL  $D_e$  (Fig. 2). The natural measurements show little sensitivity to test dose size, but the lowest test dose (2.5%) produced significant inter-aliquot scatter. The dose recovery experiment suggests underestimation at test doses  $> 30\%$  of  $D_e$ , with a relatively low test dose (8%) giving the best dose recovery ( $0.98 \pm 0.01$ ;  $n = 3$ ). For both the natural and dose recovery measurements, the DRCs behave as per Silver Lake, with faster

saturation for the lowest test doses ( $D_0$  of  $100 \pm 3$  Gy for the 2.5% test dose vs.  $306 \pm 52$  Gy for 48% test dose) and indistinguishable DRCs for moderate (23%) and high (65%) test doses (Fig. S3). Despite this, lower test doses produced the best dose recovery, with a tendency for lower  $\ln/T_n$  ratios rather than a changing DRC at high test doses. The latter was not observed in the natural  $D_e$  measurements.

#### 4.7. Fading rates

The 50 °C IR fading rates are  $10.6 \pm 2.0\%$  and  $9.4 \pm 1.0\%$  per decade for HL14-1 and 14-2, resulting in large uncertainties with fading-correction. The pIRIR<sub>225</sub> fading rates are comparable to Silver Lake ( $2.0 \pm 0.4$  and  $2.4 \pm 0.2\%$ ), but the MET-PIR plateau for HL14-2 more unambiguously implies a need for fading correction (Fig. 3), with the 250 °C age of  $35.4 \pm 2.5$  ka within uncertainties of several of Garcia et al.'s (2014) radiocarbon ages (Fig. S6). The pIRIR<sub>290</sub> ages are comparable to this ( $33.4 \pm 1.9$  ka and  $37.3 \pm 2.3$  ka for HL14-1 and HL14-2 respectively; Fig. 3 and S6), although the pIRIR<sub>290</sub> dose recovery results for HL14-1 ( $1.07 \pm 0.05$   $n = 2$ ) (natural dose plus a 66.5 Gy dose) hint at potential overestimation. The fading-corrected pIRIR<sub>225</sub> ages for both HL14-1 and HL14-2 are  $29.0 \pm 1.9$  ka and  $29.9 \pm 2.0$  ka, placing them good agreement with Meek's (1999) radiocarbon ages, but still somewhat lower than Garcia et al.'s (2014) radiocarbon ages (Fig. S6) and most of their IRSL ages.

#### 4.8. Single grain analysis

28 of 96 grains from HL14-1 produced acceptable luminescence characteristics. The distribution of grain brightness (Fig. S5) is skewed ( $\sim 20\%$  of grains account for 50% of the light sum) and the equivalent dose distribution is over-dispersed ( $29 \pm 4\%$ , with 3.3% added to individual uncertainties). A cluster of lower  $D_e$  grains are also insensitive (Fig. 4), but are not associated with higher fading rates (Fig. S7). The CAM  $D_e$  using all grains is  $94.0 \pm 6.2$  Gy, but increases to  $112.4 \pm 6.1$  Gy when the brightest 50% are used ( $n = 14$ ), and the  $D_e$  distribution becomes less dispersed ( $OD 16 \pm 4\%$ ). The (fading

uncorrected) age using the brightest grains ( $26.8 \pm 1.9$  ka) is indistinguishable from the single aliquot age. Individual grain fading rates range from 11% to –6% per decade. Although the uncertainties are large, the mean is comparable to the single aliquot analysis (2.8% per decade).

## 5. Discussion

Comparison with independent dating is limited by several factors. For Harper Lake there are inconsistencies between the radiocarbon chronologies of Meek (1999) and Garcia et al. (2014). Garcia et al.'s (2014) preferred site age was older still at 40–45 ka (citing palaeosol ages in the down-catchment Lake Manix Basin (Reheis et al., 2012) and noting that post-depositional contamination would tend to make radiocarbon ages too young; see also Reheis et al., 2015). However, the precise reasons remain unclear, and the spread of ages makes interpretation difficult (Reheis et al., 2015). The impact of the hard-water effect on radiocarbon ages was suggested by Owen et al. (2007) to be < 150 years, although Berger and Meek (1992) reported offsets up to 450 years. For luminescence ages there are also potential offsets from water content estimation. Using water contents at saturation (~35%) or akin to the modern values (2%) results in pIRIR<sub>225</sub> ages of 8.5–9.6 ka for SL14-1/LFA8 and 23.8–27.8 ka for HL14-1. Given these are extreme values, it is unlikely that this alone accounts for any differences (for Harper Lake particularly).

Nonetheless, the pIRIR<sub>225</sub> ages show substantially better agreement with the independent dating than the 50 °C IR and quartz OSL ages (Table 1; Figs. S4 and S6). The single aliquot pIRIR<sub>225</sub> data are highly reproducible and an absence of overestimation at either site implies incomplete bleaching is not an issue, despite the lacustrine contexts. Lower 50 °C IR ages reflect a need for anomalous fading correction (noting the inter-site variability) (Table 1), while quartz underestimation reflects low sensitivity and (probably) a significant contaminating non-quartz signal (Fig. S2). At Harper Lake a small proportion of quartz aliquots produce ages in better agreement with the pIRIR<sub>225</sub> ages with early background signal subtraction or if the fast ratio (Durcan and Duller, 2011) is employed as screening methods (Hay, 2018). The aliquot rejection rate is high and the ages are still lower than the fading-corrected pIRIR<sub>225</sub> ages, and much of the independent chronology. Quartz ages not employing such rigorous screening (at least, quartz of local origin) should be considered with care.

Limited sensitivity of the natural  $D_e$  to test dose size is observed for test doses between 5 and ~56% of  $D_e$  at Harper Lake and between 5 and 30% at Silver Lake. There is a clear impact on the DRC for very small test doses (Fig. S3), but this does not result in consistently higher or lower  $D_e$  estimates (note the scatter for the low test doses for the Harper Lake natural signal; Lui et al., 2016). At Harper Lake the dose recovery data appear more sensitive to test dose than the natural  $D_e$  data (Yi et al., 2016). There is a significant correlation between  $L_x$  background and the  $T_x$  initial signals for all test dose analyses (Colarossi et al., 2018), and the slope of this relationship increases at higher test doses. There is a tendency towards poorer dose recovery at high test doses at Harper Lake. This is due to a lower  $\ln/T_n$  (Fig. S3), which is also seen in natural  $D_e$  data at Silver Lake. The reason(s) for this is(are) not clear, but it implies an effect on the initial natural dose/test dose measurement.

At Silver Lake the fading-uncorrected pIRIR<sub>225</sub> ages are close (SL14-2) or fall below (14-1) the radiocarbon age ranges. For SL14-1 especially this implies fading correction (2.1% per decade) is necessary (Table 1). The MET-PIR data do not unambiguously support this however, although a small increase in the first stimulation temperature does increase the sample age. At Harper Lake most results from the independent dating and the MET-PIR/pIRIR<sub>290</sub> data more strongly indicate that fading correction of the pIRIR<sub>225</sub> ages (2.0–2.4% per decade) is required. The MET-PIR plateau (200–250 °C) and pIRIR<sub>290</sub> data fall

within the lower range of Garcia et al.'s (2014) radiocarbon dates. The fine-grain MAAD IRSL ages show less consistency than our coarse-grain pIRIR<sub>225</sub> and pIRIR<sub>290</sub> ages (samples ALG-HL-OSL2 vs ALG-HL-OSL3 in Garcia et al., 2014), which perhaps reflects uncertainty imparted when correcting the former for the high 50 °C IR fading rates at this site (Table 1), which was also based on fading analysis of a single sample.

The preliminary single grain data indicate (from dose recovery data) rather low “intrinsic” over-dispersion (using the IR LED), but this requires further investigation (c.f. Rhodes, 2015). The limited number of grains should be kept in mind. There is variability in both the signal contribution from individual grains (Fig. S5) and in the presence of a “declining baseline” (i.e. systematically lower  $D_e$ s for the dimmest grains; Rhodes, 2015, Fig. 4; Fig. S7). At Harper Lake using the brightest grains reduces OD and moves the resulting age closer to the independent dating (Lamothe et al., 2012), but the result is still within uncertainties of the age obtained using all the grains. Such a relationship is not seen at Silver Lake. At Harper Lake the insensitive, lower  $D_e$  grains do not have higher fading rates (Fig. S7). The internal K/Rb contents of the grains were not assessed, but some studies suggest that K content may not be strongly associated with grain sensitivity (Smedley et al., 2012) or fading rate (Trauerstein et al., 2014).

## 6. Conclusions

The pIRIR<sub>225</sub> protocol shows significantly better agreement with independent dating than the 50 °C IR and quartz OSL ages in the Mojave region studied. Quartz consistently and significantly underestimates expected sample ages. At both sites the pIRIR<sub>225</sub> ages show improved agreement with the independent dating after fading correction, with the MET-PIR and pIRIR<sub>290</sub> results showing even better agreement with independent  $^{14}\text{C}$  ages (of Garcia et al., 2014) at Harper Lake. However, at the younger Silver Lake site the SL14-1 MET-PIR and pIRIR<sub>290</sub> ages are identical to the uncorrected pIRIR<sub>225</sub> age. The pIRIR<sub>225</sub> measurements show limited sensitivity to test dose size at Harper Lake, but at both sites the DRCs saturate faster for very low test doses and underestimate for high test doses at Silver Lake. The latter is not observed at Harper Lake where the dose recovery data seem to be more sensitive to test dose size than the natural  $D_e$  measurements. Contrasting single grain behaviors are also observed, notably in the presence of less sensitive, lower  $D_e$  grains at Harper Lake. This mirrors some previous work in suggesting, at least for some sites, that the brightest K-feldspar grains may provide a better estimate of burial dose.

## Acknowledgments

ASH was supported by NERC studentship 1358108. Rob Fulton, Jason Wallace and Simon Benson are thanked for logistical support. An anonymous reviewer is thanked for very constructive comments.

## Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.quageo.2018.05.006>.

## References

- Auclair, M., Lamothe, M., Huot, S., 2003. Measurement of anomalous fading for feldspar IRSL using SAR. *Radiat. Meas.* 37, 487–492.
- Bateman, M.D., Bryant, R.G., Foster, I.D.L., Livingstone, I., Parsons, A.J., 2012. On the formation of sand ramps: a case study from the Mojave Desert. *Geomorphology* 161–162, 93–109.
- Berger, R., Meek, N., 1992. Radiocarbon dating of *Anodonta* in the Mojave River basin. *Radiocarbon* 34, 578–584.
- Brown, N.D., Rhodes, E.J., Antinao, E.V., McDonald, E.V., 2015. Single grain post-IR IRSL signals of K feldspar of alluvial fan deposits in Baja California Sur, Mexico. *Quat. Int.* 362, 132–138.
- Buylaert, J.-P., Jain, M., Murray, A.S., Thomsen, K.J., Theil, C., Sohbaty, R., 2012. A robust feldspar luminescence dating method for Middle and Late Pleistocene

- sediments. *Boreas* 41, 435–451.
- Colarossi, D., Duller, G.A.T., Roberts, H.M., 2018. Exploring the behaviour of luminescence signals from feldspars: implications for the single aliquot regenerative dose protocol. *Radiat. Meas.* 109, 34–44.
- Clarke, M.L., Richardson, C.A., Rendell, H.M., 1995. Luminescence dating of Mojave Desert sands. *Quat. Sci. Rev.* 14, 783–789.
- Durcan, J.A., Duller, G.A.T., 2011. The fast ratio: a rapid measure for testing the dominance of the fast component in the initial OSL signal from quartz. *Radiat. Meas.* 46, 1065–1107.
- Enzel, Y., Wells, S.G., Lancaster, N., 2003. Late Pleistocene lakes along the Mojave River, southeast California. *Geol. Soc. Am. Spec. Pap.* 368, 61–78.
- Fuchs, M., Dietze, M., Al-Qudah, K., Lomax, J., 2015. Dating desert pavements – first results from a challenging environmental archive. *Quat. Geochronol.* 30, 342–349.
- Garcia, A.L., Knott, J.R., Mahan, S.A., Bright, J., 2014. Geochronology and paleoenvironment of pluvial Harper Lake, Mojave Desert, California, USA. *Quat. Res.* 81, 305–317.
- Hay, A.S., 2018. The Influence of Complex Topography on Aeolian Sediment Accumulation and Preservation: an Investigation of Morphology and Process History. Unpublished PhD thesis. University of Leicester.
- Huntley, D.J., Lamothe, M., 2001. The ubiquity of anomalous fading in K-feldspars and the measurement and correction for it in optical dating. *Can. J. Earth Sci.* 38, 1093–1106.
- Kars, R.H., Reimann, T., Wallinga, J., 2014. Are feldspar SAR protocols appropriate for post-IR IRSL dating? *Quat. Geochronol.* 22, 126–136.
- Kreutzer, S., Schmidt, C., Fuchs, M.C., Dietze, M., Fisher, M., Fuchs, M., 2012. Introducing an R package for luminescence dating analysis. *Ancient TL* 30, 1–7.
- Lamothe, M., Barré, M., Huot, S., Ouimet, S., 2012. Natural luminescence and anomalous fading in K-feldspar. *Radiat. Meas.* 47, 682–687.
- Lancaster, N., Tchakerian, V.P., 1996. Geomorphology and sediments of sand ramps in the Mojave. *Geomorphology* 17, 151–165.
- Lawson, M.J., Roder, B.J., Stang, D.M., Rhodes, E.J., 2012. OSL and IRSL characteristics of quartz and feldspar from southern California, USA. *Radiat. Meas.* 47, 830–836.
- Li, B., Li, S.-H., 2011. Luminescence dating of K-feldspar from sediments: a protocol without anomalous fading correction. *Quat. Geochronol.* 6, 468–479.
- Li, B., Jacobs, Z., Roberts, R.G., Li, S.-H., 2014. Review and assessment of the potential of post-IR IRSL dating methods to circumvent the problem of anomalous fading in feldspar luminescence. *Geochronometria* 41, 178–201.
- Lui, J., Murray, A.S., Sohbati, R., Jain, M., 2016. The effect of test dose and first IR stimulation temperature on post-IR IRSL measurements of rock slices. *Geochronometria* 43, 179–187.
- McGuire, C., Rhodes, E.J., 2015. Determining fluvial sediment virtual velocity on the Mojave River using K-feldspar IRSL: initial assessment. *Quat. Int.* 362, 124–131.
- Meek, N., 1999. New Discoveries About the Late Wisconsinan History of the Late Mojave River System. San Bernadino County Museum Association Quarterly 46, pp. 113–117.
- Murray, A.S., Wintle, A.G., 2000. Luminescence dating of quartz using an improved single-aliquot regenerative-does protocol. *Radiat. Meas.* 32, 57–73.
- Ore, H.T., Warren, C.N., 1978. Late pleistocene-early holocene geomorphic history of Lake Mojave, California. *Geol. Soc. Am. Bull.* 82, 2553–2562.
- Owen, L.A., Bright, J., Finkel, R.C., Jaiswal, M.K., Kaufman, D.S., Mahan, S., Radtke, U., Schneider, J.S., Sharp, W., Singhvi, A.K., Warren, C., 2007. Numerical dating of a Late Quaternary spit-shoreline complex at the northern end of Silver Lake playa, Mojave Desert, California: a comparison of the applicability of radiocarbon, luminescence, terrestrial cosmogenic nuclide, electron spin resonance, U-series and amino acid racemization methods. *Quat. Int.* 166, 87–110.
- Reheis, M.C., Bright, J., Lund, S.P., Miller, D.M., Skipp, G., Fleck, R.J., 2012. A half-million-year record of paleoclimate from the Lake Manix core, Mojave Desert, California. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 365, 11–37.
- Reheis, M.C., Miller, D.M., McGeehin, J.P., Redwine, J.R., Oviatt, C.G., Bright, J., 2015. Directly dated MIS 3 lake-level record from Lake Manix, Mojave desert, California, USA. *Quat. Res.* 83, 187–203.
- Rendell, H.M., Sheffer, N.L., 1996. Luminescence dating of sand ramps in the eastern Mojave Desert. *Geomorphology* 17, 187–197.
- Rhodes, E.J., 2015. Dating sediments using potassium feldspar single-grain IRSL: initial methodological considerations. *Quat. Int.* 362, 14–22.
- Roder, B., Lawson, M., Rhodes, E.J., Dolan, J., McAuliffe, L., McGill, S., 2012. Assessing the potential of luminescence dating for fault slip rate studies on the Garlock fault, Mojave Desert, California, USA. *Quat. Geochronol.* 10, 285–290.
- Smedley, R.K., Duller, G.A.T., Pearce, N.J.G., Roberts, H.M., 2012. Determining the K-content of single-grains of feldspar for luminescence dating. *Radiat. Meas.* 47, 790–796.
- Thomsen, K.J., Murray, A.S., Jain, M., Bøtter-Jensen, L., 2008. Laboratory fading rates of various luminescence signals from feldspar-rich sediment extracts. *Radiat. Meas.* 43, 1474–1486.
- Trauerstein, M., Lowick, S.E., Preusser, F., Schlunegger, F., 2014. Small aliquot and single grain IRSL and post-IR IRSL dating of fluvial and alluvial sediments from the Pativilca Valley, Peru. *Quat. Geochronol.* 22, 163–174.
- Vermeesch, P., 2009. RadialPlotter: a Java application for fission track, luminescence and other radial plots. *Radiat. Meas.* 44, 409–410.
- Wells, S.G., Brown, W.J., Enzel, Y., Anderson, R.Y., McFadden, L.D., 2003. Late Quaternary geology and paleohydrology of pluvial Lake Mojave, southern California. *Geol. Soc. Am. Spec. Pap.* 358, 79–114.
- Yi, S., Buylaert, J.-P., Murray, A.S., Lu, H., Thiel, C., Zeng, L., 2016. A detailed post-IR IRSL dating study of the Niuyangzigou loess site in northeastern China. *Boreas* 45, 644–657.