



Review

Epigenetics of concussion: A systematic review

Mark R. Antrobus^{a,*}, Terun Desai^b, David Young^a, Lee Machado^a, William J. Ribbans^a,
Louis Y. El Khoury^c, Jon Brazier^d

^a Centre for Physical Activity and Life Sciences, University of Northampton, Northampton NN1 5PH, UK

^b Institute of Sport, Exercise & Health, Division of Surgery & Interventional Science, University College London, W1T 7HA, UK

^c Department of Pathology, St. Jude Children's Research Hospital, Memphis, TN, USA

^d Department of Psychology, Geography and Sport, University of Hertfordshire, Hatfield AL10 9AB, UK

ARTICLE INFO

Keywords:

Epigenetics
Concussion
Brain
miRNA
Methylation
Histone

ABSTRACT

Background: Concussion is the most common neurological disorder affecting millions of people globally each year. Identifying epigenetic mechanisms influencing concussion incidence, severity and recovery could provide diagnostic and prognostic insight into this injury.

Objectives: This systematic review aims to identify the epigenetic mechanisms underpinning concussion.

Methods: Seven electronic databases; PubMed, MEDLINE, CINAHL, Cochrane library, SPORTDiscus, Scopus and Web of Science were searched for studies that investigated the epigenetic mechanisms of concussion and its underlying neuropathology.

Results: Based on inclusion and exclusion criteria, 772 titles were independently analysed by two of the authors to a final list of 28 studies that totaled 3042 participants. We observed separate associations between sncRNAs, methylation, histone modification and concussion. Overall, 204 small non-coding RNAs were significantly dysregulated between concussed participants and controls or between concussion participants with no post-concussive symptoms and those with post-concussive symptoms. From these, 37 were reported in more than one study and 23 of these were expressed in a consistent direction with at least one further study. Ingenuity pathway analysis identified 10 miRNAs known to regulate 15 genes associated with human neurological pathologies. Two studies found significant changes in global methylation in concussed participants and one study found a decrease in H3K27Me3 in the context of DNA damage and concussion.

Conclusions: The review findings suggest that epigenetic mechanisms may play an important role in the pathophysiological mechanisms that could influence outcome, recovery, and potential long-term consequences of concussion for individuals.

Key Points

Epigenetic mechanisms may play an important role in the pathophysiological mechanisms that could influence risk, outcome, recovery, and potential long-term consequences of concussion for individuals.

A better understanding of epigenetic mechanisms that contribute to the neuropathology of concussion is essential to improve understanding and subsequent management of this form of traumatic brain injury and to provide directions for future study.

1. Introduction

Traumatic brain injury (TBI) is the most common neurological disorder affecting 50–69 million people globally each year (Dewan et al., 2018; Lefevre-Dognin et al., 2021). The severity of TBI varies from severe to mild, with the latter accounting for most cases (70–95 %) presented at hospital (Maas et al., 2022; Maas et al., 2017). In addition, incidence is underestimated as up to 50 % of concussions may go undiagnosed (Broglia et al., 2017; Meehan et al., 2013; McCrea et al., 2004). Concussion is a subgroup of TBI also termed as mild-TBI (mTBI)

Abbreviations: TBI, Traumatic brain injury; mTBI, mild-TBI; GWAS, Genome-wide-association study; SNPs, Single nucleotide polymorphisms; miRNAs, microRNAs; piRNAs, PIWI-interacting RNAs; siRNAs, small interfering RNAs; snoRNAs, small nucleolar RNAs; GCS, Glasgow Coma Score; NOS, Newcastle-Ottawa Scale; CSF, cerebral spinal fluid.

* Corresponding author.

E-mail address: mark.antrobus@northampton.ac.uk (M.R. Antrobus).

<https://doi.org/10.1016/j.gene.2024.149046>

Received 8 July 2024; Received in revised form 14 October 2024; Accepted 22 October 2024

Available online 26 October 2024

0378-1119/© 2024 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY-NC license (<http://creativecommons.org/licenses/by-nc/4.0/>).

however, within the related literature a wide variety of definitions are used which has implications for study comparison. Sustaining a concussion elicits altered brain function via complex pathophysiological processes that can have both short and long-term consequences (Barkhoudarian et al., 2016). Most individuals recover from concussion within 2–4 weeks post-concussive event, however 10 % of patients experience persisting symptoms after concussion for 1–3 months or longer (Willer and Leddy, 2006). The potential effects of sustaining a concussion may have long-lasting functional and structural changes to the frontal and temporal regions of the brain, as observed in certain individuals who sustained a concussion 20 years prior (June et al., 2020). In addition, sustaining repetitive concussions may be a risk factor for developing progressive neurodegenerative diseases such as chronic traumatic encephalopathy and tauopathies (Buckland et al., 2022;13:1107.; Nowinski et al., 2022; Mckee and Daneshvar, 2015; Russell et al., 2022; Bieniek et al., 2020).

Several environmental risk factors contribute to concussion risk such as, behaviour, rules of the sport, nutrition, quality of sleep and prior history of concussion (Ponsford et al., 2000; Abrahams et al., 2014; Iverson et al., 2017; Raikes et al., 2019; Lust et al., 2020). Additionally, genetic risk factors may affect incidence and outcomes related to concussion (Panenka et al., 2017; Antrobus et al., 2021). A recent genome-wide-association study (GWAS) identified 2 novel SNPs (*SPATA5* rs144663795 and *PLXNA4* rs117985931) associated with concussion (Kim et al., 2020). Currently the heritability of concussion is unknown. Based on other observations of substantial genetic contributions to inter-individual variability in most human traits, it is likely that a substantial genetic component applies to concussion (Antrobus et al., 2021). However, these genetic variants only partially explain the inter-individual variability of incidence and outcomes related to concussion. In addition, epigenetic mechanisms are likely to also contribute to the neuropathophysiology of concussion.

Epigenetics refers to functional changes to the genome that regulate gene expression in response to environmental stimuli, without changing the primary DNA base sequence (Waddington, 1942). The primary epigenetic mechanisms are DNA methylation, histone modifications, and the post-transcriptional regulation by small non-coding RNAs (sncRNAs, which include microRNAs (miRNAs), PIWI-interacting RNAs (piRNAs), small interfering RNAs (siRNAs), and small nucleolar RNAs (snoRNAs)). DNA methylation occurs by the addition of a methyl group to the C-5 carbon atom position of cytosine bases to form 5' methyl-cytosine which alters gene expression by recruiting proteins involved in gene repression or by inhibiting transcription factor(s) binding to DNA (Moore et al., 2013). Histone acetylation, phosphorylation, and methylation are epigenetic modifications that are associated with altered gene transcription (Bannister et al., 2011). Small non-coding RNAs are highly abundant molecules that regulate essential cellular functions with a wide variety of roles. For example, miRNAs directly control transcription by recruiting the machinery for gene silencing of gene promoter in various neurodegenerative diseases such as Alzheimer's disease and Parkinson's disease (Li et al., 2022). Similarly, piRNAs silence gene expression through transcription and post-transcriptional processes and within the brain can play a role in neuronal differentiation and development of neurodegenerative diseases (Sato et al., 2023). siRNAs regulate the expression of genes through the process of RNA interference (Dana et al., 2017) and finally snoRNAs stabilise the structure of rRNA through modifying rRNA with 2'-O-methylation and pseudouridylation of nucleotides (Stepanov et al., 2015;2015.).

To date, there is little current evidence delineating the role of epigenetic mechanisms in concussion. Therefore, we have systematically reviewed studies which have investigated the epigenetic mechanisms of concussion. A better understanding of epigenetic mechanisms that contribute to the neuropathology of concussion is essential to improve understanding and subsequent management of this form of TBI and to provide directions for future study.

2. Methods

A systematic review assessed the evidence for epigenetic mechanisms of concussion and how they contribute to the neuropathology of concussion. This systematic review was developed in accordance with the guidelines of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) (Page et al., 2020). A review protocol was registered with PROSPERO on 15th September 2022.

2.1. Search terms and databases

Keywords were selected by agreement of all authors and drafted into a PubMed search strategy. The search strategy was reviewed and revised using the CADTH Peer Review Check-list for Search Strategies according to the PRESS 2015 Guideline Statement, and then tested to ensure key studies were retrieved (McGowan et al., 2015). The PubMed search (Table 1) was adapted for all other databases searched in this review (MEDLINE, CINAHL, Cochrane library, SPORTDiscus, Scopus and Web of Science). Results were filtered to include studies in peer-reviewed journals (no date restriction) and were limited to English language articles only. The search was conducted on 20th May 2024.

Table 1. Schematic to represent 2-level search strategy.

2.2. Study selection criteria and data extraction

Studies were included if they (i) Were primary research; (ii) Included concussion participants (those of any age or sex, diagnosed with a concussion, occurring from any sport, activity, combat, accident, or life event) diagnosed according to a form of diagnosis e.g. GCS \geq 13 or self-reported history of concussions; and (iii) Contained epigenetic mechanism (DNA methylation, histone modification and sncRNAs) study in humans only. Studies were excluded if they contained the following: (i) Participants with a moderate or severe TBI, or mixed population of mild, moderate or severe traumatic brain injury or penetrating head injury; (ii) Intervention (use of drugs alone or with physical exercise, intervention not clear); (iii) Animal based studies; and (iv) design (reviews, letters to the editor, case reports, commentaries, editorials, expert opinion, theses or dissertations). Initially, citations were independently screened by pairs of authors according to the inclusion and exclusion criteria. When required, a third reviewer adjudicated disagreements between authors. Study title and abstracts were screened for suitability and full text articles were retrieved. Data was extracted by one author and reviewed by a second author to ensure accuracy and completeness.

2.3. Assessment of risk of bias

For each included study the risk of bias was independently rated by two authors using the Newcastle-Ottawa Scale (NOS) in line with previous studies (Bannister et al., 2011; Li et al., 2022), with a third reviewer adjudicating disagreements between authors. A score of nine stars deemed a study to be of good quality and at low risk of bias; studies

Table 1
Schematic to represent 2-level search strategy.

Operator	Search terms
	#1 epigen* OR DNA methylation OR S-Adenosylmethionine OR hypermethylation OR hypomethylation OR CpG Islands OR Histone OR acetylation OR demethylation OR methylation OR phosphorylation OR ubiquitination OR modification OR adenosylmethionine OR CpG OR microRNA* OR miRNA OR mirna biomarker* OR non-coding RNA* OR gene modification OR gene expression
AND	#2 mild traumatic brain injur* OR MTBI OR concussion* OR brain concussion* OR post-concussion* OR sports-related concussion*
NOT	Animal* NOT rat* NOT mouse NOT mice NOT murine NOT dog* NOT canine NOT cat* NOT feline

that scored eight or seven stars were deemed at medium risk of bias; while those that scored below seven were at high risk of bias.

2.4. Ingenuity pathway analysis

Qiagen's Ingenuity Pathways Analysis (IPA, v23.0) was used to identify pathways based on post-concussion miRNA expression profiles. To reduce the number of miRNA-mRNA interactions to a workable number, we employed filters for miRNA confidence (experimentally observed), disease (neurological diseases), pathways (neuronal system), species (human), and tissue (nervous system).

3. Results

3.1. Literature search Results

The search of the databases identified 735 articles (Fig. 1). After duplicates were removed and each article's reference list was hand searched for other relevant articles, 575 records were screened and a total of 28 were included in this review. The NOS scores of the case-control studies ranged from 6 to 8, with a mean score of 7.6. The NOS scores of the cohort studies ranged from 7 to 8, with a mean score of 7.6. The overall mean NOS score across all studies was 7.6 (Table 2). Each of the 28 studies were critically reviewed by at least two investigators using a standard review form.

3.2. Characteristics of the included studies

The characteristics and key findings are summarised including; participant characteristics, concussion diagnosis, epigenetic mechanism and main findings (Table 3). Of the 28 included studies 16 were case-control studies and the remaining 12 were cohort studies. The total

number of participants across all studies was 3042, ranging from 18 to 538 participant sample sizes, with a median sample size of 55. The mean age range of participants in the selected articles was 12.8–54.6 years, seven studies (25 %) included children (<18 years). Twenty-two studies (79 %) included both male and female participants and only 6 (21 %) studies included only male participants. Only one study (4 %) had white only participants. Fifteen studies (54 %) used a participant sample of mixed ethnic groups, and twelve studies (43 %) did not report the ethnicity of participants.

Five studies included military personnel, four studies included only participants under the age of 18 years old, two studies focused on professional rugby union athletes, one study included Australian Football League athletes, one study contained professional contact athletes from a variety of sports, four studies included professional American football athletes, three studies contained collegiate athletes from a mix of sports and finally eight studies focused on the general population (some of which included participants under the age of 18 years old). Most studies used whole blood collection (17), nine studies utilised saliva, one study collected cerebral spinal fluid (CSF) and one collected human brain tissue. A variety of methods were utilised to measure the epigenetic response of concussion with six studies using miRNA from saliva, six studies used miRNA from serum, four studies used miRNA from plasma, two studies utilised miRNA from plasma and extracellular vesicles, two studies used miRNA from exosomal plasma, two studies measured DNA methylation at 850,000 CpG sites, one study measured global DNA methylation (5-mC%), one study used miRNA from CSF, one study measured histone modification from brain tissue, with the remaining three studies measuring small nuclear RNA/miRNA and sncRNA from blood and saliva, respectively.




Fig. 1. PRISMA flow diagram.

Table 2
Quality assessment of studies.

Study	Study design	Newcastle-Ottawa score			Total
		Selection	Comparability	Outcome/exposure	
Devoto et al. (Devoto et al., 2022)	CC	****	**	**	8
Shultz et al. (Shultz et al., 2022)	CH	****	*	***	8
Di Pietro et al. (Di Pietro V, O'Halloran P, Watson CN, Begum G, Acharjee A, Yakoub KM, et al. Unique diagnostic signatures of concussion in the saliva of male athletes: the Study of Concussion in Rugby Union through MicroRNAs (SCRUM). Br J Sports Med [Internet]. Br J Sports Med; 2021)	CH	***	*	***	7
Duan et al. (Duan et al., 2021)	CC	****	**	**	8
Fedorchak et al. (Fedorchak et al., 2021)	CC	***	*	***	7
Guedes et al. (Guedes et al., 2021)	CH	***	**	***	8
Hicks et al. (Hicks et al., 2021)	CC	****	**	**	8
Lusardi et al. (Lusardi et al., 2021)	CC	****	**	**	8
Papa et al. (Papa et al., 2021)	CH	***	*	***	7
Vorn et al. (Vorn et al., 2021)	CC	****	*	*	6
Bahado-Singh et al. (Bahado-Singh et al., 2020)	CC	****	*	**	7
Ghai et al. (Ghai et al., 2020)	CC	****	**	**	8
Hicks et al. (Hicks et al., 2020)	CC	****	*	**	7
Lee et al. (Lee et al., 2020)	CC	****	**	**	8
Polito et al. (Polito et al., 2020)	CC	****	**	**	8
Davies et al. (Davies et al., 2019)	CC	****	**	**	8
Papa et al. (Papa et al., 2019)	CH	***	*	**	6
Schwab et al. (Schwab et al., 2019)	CC	****	**	**	8
Svingos et al. (Svingos et al., 2019)	CH	***	**	***	8
Di Pietro et al. (Di Pietro et al., 2018)	CC	****	*	**	8
Johnson et al. (Johnson et al., 2018)	CH	***	**	***	8
Pasinetti et al. (Pasinetti et al., 2012)	CC	****	*	**	7
Hicks et al. (Hicks et al., 2020)	CC	****	**	**	8
Miller et al. (Miller et al., 2022)	CH	***	**	***	8
Mitra et al. (Mitra et al., 2022)	CH	***	**	***	8
Mitra et al. (Mitra et al., 2023)	CH	***	**	***	8
Tas et al. (Tas et al., 2020)	CH	***	**	***	8
Wyczechowska et al. (Wyczechowska et al., 2023)	CH	***	**	***	8

CC, case-control; CH, cohort.

3.3. Epigenetic mechanisms associated with concussion

Across the 28 studies identified, 24 of these investigated sncRNAs. There was a large variation in the expression of sncRNAs in concussed patients between studies (Table 3). Specifically, there was a total of 204 sncRNAs that were significantly up- or down-regulated between concussed patients and controls or between concussion patients with no post-concussive symptoms and those with post-concussive symptoms (Table 3). From these, 37 were reported in more than one study and 23 of these miRNAs were reported to be differentially expressed in a consistent direction with at least one further study. Table 4 summarises the miRNAs identified in multiple studies. Twenty-two miRNAs were identified in two or more studies which could have potential diagnostic roles in the management of concussion (Fig. 2). Thirty-four miRNAs were identified in two or more studies which could have potential prognostic roles in the management of concussion (Fig. 2).

For the remaining four studies, three investigated DNA methylation. Lee et al. identified higher blood global methylation ratio (5-mC%, percentage of methylated cytosines in total DNA) in concussed cases (4.42 ± 0.43) than in controls (3.93 ± 0.03) (Lee et al., 2020). Bahado-Singh et al. found 449 CpG sites were significantly methylated with some methylation changes > 10 % in concussed youth compared to youth controls (Bahado-Singh et al., 2020). However, Duan et al. did not identify any significant global methylation changes between a concussion group and controls (Duan et al., 2021). Finally, Schwab et al. examined histone modification in human brain tissue, finding cases with a history of concussion had a decrease of H3K27Me3 in the context of DNA damage (Schwab et al., 2019).

To investigate the potential contribution of concussion-associated candidate miRNAs, ingenuity pathway analysis (IPA) was performed employing the 37 miRNAs to identify putative miRNA targets relevant to the nervous system and neuronal disease (Fig. 2). This resulted in identification of 10 miRNAs known to regulate 15 genes (*ESR1*, *NR3C2*,

PGR, *CREB1*, *PDPK1*, *PRKCA*, *AP2A1*, *STX1A*, *KRAS*, *ADCY6*, *ERBB4*, *GRIA2*, *SLC6A4*, *GNAI2*, *RAC1*) that play a role in human neurological pathologies (Fig. 3). Based on the post-concussion directions of change reported in Table 4, we decided to build a consensus prediction model, i. e. a miRNA is said to be upregulated (coloured in red, let-7a-5p, miR-142-3p, miR-135b-5p, and miR-372-3p) if it is upregulated in a majority of papers. The opposite is true for downregulated miRNAs (coloured in green, miR-30e-5p, miR-181a-5p, miR-27a-3p, and miR199a-3p). Where a consensus cannot be reached because there is an even number of inconsistent findings, the miRNA is not coloured (miR-182-5p and miR-20a-5p). The phenotypes predicted by IPA based on the consensus approach, show an activation of brain concussion, neuronal cell death, long-term synaptic depression of brain cells, and long-term potentiation of brain. Conversely, there is an inhibition of long-term synaptic depression of neurons, inhibition of proliferation of neuronal cells, and inhibition of neuron density.

4. Discussion

This systematic review identified 28 studies suitable for establishing our present understanding of the epigenetic mechanisms associated with concussion (Table 2 and 3). The methods adopted across these studies included case studies, cohort studies, clinical assessments, self-reporting of concussion, neuroimaging and autopsy. Separate associations between sncRNAs, methylation, histone modification and concussion were observed within the selected studies. The earliest published study in this review dates back only to 2012 suggesting that there is a relatively nascent but growing literature on epigenetic mechanisms that may play an important role in the pathophysiological mechanisms that influence outcome, recovery and potential long-term consequences following concussion for individuals.

Table 3
Characteristics and key findings of epigenetics of concussion studies.

Study	Participants N	Sex	Ethnicity	Age (mean) [Years]	Concussion Diagnosis	Epigenetic mechanism	Tissue type	Main Findings
Devoto et al. (Devoto et al., 2022)	152 military service members and veterans with a history of concussion, 35 controls without history of concussion.	163 males, 24 females	Not reported	38.0	Modified OSU mTBI-ID, VCU rCDI (mean time since last concussion ranged between 8.0–12.2 years for all groups).	miRNAs (exosomal plasma)	Whole blood	23 miRNAs associated with pathways involved in neuronal function, chronic inflammatory processes and neuronal death were upregulated in the blast-related concussion group compared to controls (hsa-let7e-5p, hsa-miR-1233-3p, hsa-miR-1268b, hsa-miR-1304-3p, hsa-miR-18a-5p, hsa-miR-197-5p, hsa-miR-204-5p, hsa-miR-3190-3p, hsa-miR-324-3p, hsa-miR-326, hsa-miR-361-3p, hsa-miR-3615, hsa-miR-372-3p, hsa-miR-376b-3p, hsa-miR-4792, hsa-miR-5001-5p, hsa-miR-5010-3p, hsa-miR-516a-5p, hsa-miR-567, hsa-miR-615-5p, hsa-miR-619-3p, hsa-miR-631, hsa-miR-767-5p) and 1 downregulated (hsa-miR-139-5p).
Shultz et al. (Shultz et al., 2022)	28 AFL athletes with a history of concussion, 27 AFL athlete controls with no history of concussion.	39 males, 16 females	Not reported	24.0	GCS (samples were collected pre-season, 2-, 6-, and 13-days post-concussion).	miRNA (plasma)	Whole blood	miR-221-3p levels were decreased at 6- and 13-days post-concussion, and miR-27a-3p levels were decreased at 6-days post-concussion, when compared to baseline. Both miR-27a and miR-221-3p levels were inversely correlated with concussion symptom severity. miR-221 plays a role in neuronal cell differentiation vascular homeostasis, miR-27a has been related to regulation of neuroinflammation.
Di Pietro et al. (Di Pietro V, O'Halloran P, Watson CN, Begum G, Acharjee A, Yakoub KM, et al. Unique diagnostic signatures of concussion in the saliva of male athletes: the Study of Concussion in Rugby Union through MicroRNAs (SCRUM). Br J Sports Med [Internet]. Br J Sports Med; 2021)	106 rugby athletes with concussion, 50 rugby athletes for who concussion was ruled out, 102 non-injured control rugby athletes.	258 males	Mixed ethnicities	26.7	HIA and SCAT-5 (samples were collected in-match, post-match and 36–48 h post-match).	miRNA (salivary)	Saliva	14 miRNAs associated with neuroinflammation and roles in AD and depression, could differentiate concussed athletes from all other athlete groups, immediately after a match (let-7a-5p, miR-143-3p, miR-103a-3p, miR-34b-3p, RNU6-7, RNU6-45, snora57, snoU13.120, tRNA18Arg-CCT, U6-168, U6-428, U6-1249, Uco22c1, YRNA-255).
Fedorchak et al. (Fedorchak et al., 2021)	32 youth participants with a persistent post-concussion symptoms, 80 youth control participants with non-persistent post-concussion symptoms history of concussion	63 males, 49 females	Mixed ethnicities	15.9	GCS scores (within 24 h post-injury)	ncRNA (salivary)	Saliva	16 ncRNAs demonstrated prognostic utility for persisting symptoms after concussion (7 miRNAs; hsa-miR-486-5p, hsa-miR-1246, hsa-miR-92b-3p, hsa-miR-203a-5p, hsa-miR-148a-5p, hsa-miR-100-5p, hsa-miR-148a-5p; 8 wiRNAs; wiRNA-48, wiRNA-176, wiRNA-1500, wiRNA-9924,

(continued on next page)

Table 3 (continued)

Study	Participants N	Sex	Ethnicity	Age (mean) [Years]	Concussion Diagnosis	Epigenetic mechanism	Tissue type	Main Findings
Guedes et al. (Guedes et al., 2021)	71 military service members and veteran with a history of concussion and no PTSD, 45 military service members and veteran with a history of concussion and PTSD, 28 controls with no history of concussion and no PTSD.	125 males, 19 females	Not reported	37.0	GCS scores, VCU rCDI (mean time since last concussion was 9 years).	miRNA (plasma and extracellular vesicles)	Whole blood	wiRNA-7971, wiRNA-1385, wiRNA-7876; 1 snoRNA; SNORD81). 12 miRNAs associated with neurodegenerative and inflammatory processes were differentially regulated between groups; concussion history and no PTSD group vs controls (hsa-miR-139-5p, hsa-miR-204-5p, hsa-miR-372-3p, hsa-miR-509-3-5p, hsa-miR-615-5p, hsa-miR-1277-3p); concussion history and PTSD group vs controls (hsa-miR-3190-3p, hsa-miR-615-5p, hsa-miR-1185-1-3p, hsa-miR-3196, hsa-miR-372-3p, hsa-miR-139-5p).
Hicks et al. (Hicks et al., 2021)	75 participants with concussion, 97 control participants with no history of concussion.	107 males, 65 females	Mixed ethnicities	20.4	GCS scores (within 24 h post-injury)	miRNA (salivary)	Saliva	miR-27a-5p was the most accurate miRNA for differentiating concussed and non-concussed participants. A logistic regression model using a ratio of miR-27a-5p and miR-30a-3p provided the highest accuracy for differentiating concussion status.
Lusardi et al. (Lusardi et al., 2021)	45 male military veterans with a history of concussion, 18 male military control veterans without a history of concussion, 52 male civilian controls a without history of concussion.	115 males	Mixed ethnicities	33.1	Clinical history via semi-structured interview (mean time between sample collection and the last concussion experienced was 5 years).	miRNA (CSF)	CSF	10 miRNAs were identified with differential expression between veterans with history of concussion and civilian controls (miR-502-3p, miR-362-3p, miR-191-5p, miR-197-3p, miR-30c-5p, miR-140-5p, miR-30b-5p, miR-20a-5p, miR-548a-3p, miR-20b-5p). 8 miRNAs were identified with differential expression between veterans with no history of concussion and veterans with a history of concussion (miR-191-5p, miR-152-3p, miR-132-3p, miR-362-5p, miR-548c-3p, miR-125a-5p, miR-130a-3p, miR-411-5p).
Papa et al. (Papa et al., 2021)	14 collegiate American Football athletes with a history of concussion, 30 non-athlete controls with no history of concussion.	29 males, 15 females	Not reported	26.5	Heads Up Centres for Disease Control's Acute Concussion Evaluation tools (samples were collected within 24 h of injury).	miRNA (serum)	Whole blood	3 miRNAs that play a role in neurocognitive functioning (miR-505, miR-30d, miR-92a) were associated with lower fractional anisotropy values of white matter dorsal (posterior) cervical spinal tracts in athletes.
Vorn et al. (Vorn et al., 2021)	29 participants with a history of concussion, 11 control participants with no history of concussion.	19 males, 21 females	Mixed ethnicities	24.4	Self-reported history of concussion diagnosed using ACRM guidelines (mean time between sample collection and the last concussion experienced was 4 years).	miRNAs (exosomal plasma)	Whole blood	25 significantly dysregulated miRNAs associated with neurological disease, organismal injury and psychological disorders were identified in concussion group compared to controls (upregulated; hsa-miR-520e, hsa-miR-499b-3p, hsa-miR-520b, hsa-miR-4488) (downregulated; hsa-miR-

(continued on next page)

Table 3 (continued)

Study	Participants N	Sex	Ethnicity	Age (mean) [Years]	Concussion Diagnosis	Epigenetic mechanism	Tissue type	Main Findings
								625-5p, hsa-miR-421, hsa-miR-664a-3p, hsa-miR-28-3p, hsa-miR-125a-5p, hsa-miR-222-3p, hsa-miR-140-5p, hsa-miR-98-5p, hsa-miR-148a-3p, hsa-miR-423-5p, hsa-miR-107, hsa-miR-181a-5p, hsa-miR-374a-5p, hsa-miR-340-5, hsa-miR-29b-3p, hsa-miR-191-5p, hsa-miR-199a-3p, hsa-miR-126-3p, hsa-miR-23a-3p, hsa-miR-142-3p, hsa-miR-223-3p).
Ghai et al. (Ghai et al., 2020)	27 military veterans with a history of concussion, 11 military control veterans without a history of concussion, 31 civilian controls without a history of concussion.	69 males	Mixed ethnicities	31.9	Clinical examinations and semi-structured interviews (time lapse since the last concussion and sample collection was 1.6–7.7 years).	miRNA (plasma and extracellular vesicles)	Whole blood	32 miRNAs associated with pathways involved in neuronal function, vascular remodelling, blood–brain barrier integrity, and neuroinflammation in plasma were significantly changed in the chronic concussion veterans group compared with control groups (let-7f-1-5p, miR-103a-1-3p, miR-103b-1-5p, miR-106a-5p, miR-106b-5p, miR-1246-5p, miR-132-5p, miR-142-3p, miR-144-3p, miR-144-5p, miR-148a-5p, miR-15a-5p, miR-15b-5p, miR-16-1-5p, miR-17-5p, miR-182-5p, miR-183-5p, miR-184-3p, miR-18a-5p, miR-19a-3p, miR-20a-5p, miR-20b-5p, miR-223-3p, miR-2355-3p, miR-32-5p, miR-3613-5p, miR-374b-5p, miR-411-5p, miR-4440-3p, miR-484-5p, miR-490-3p, miR-96-5p).
Hicks et al. (Hicks et al., 2020)	13 American Football athletes with history of concussion, 18 aged-match controls, 310 controls with mixed history of concussion	239 males, 102 females	Mixed ethnicities	47.0	Diagnosis method of concussion history not reported.	miRNA (salivary)	Saliva	3 miRNAs (miR-28-3p, miR-339-3p, and miR-361-5p) differentiated between the older former professional athletes and younger individuals with prior concussion. In the younger group the same 3 miRNAs differentiated between participants with a history of concussion and those without.
Polito et al. (Polito et al., 2020)	20 participants with a history of concussion, 10 matched 'normal' controls.	15 males, 15 females	Not reported	37.8	GCS (samples were collected at 0 h, 24 h and 48 h post-concussion).	miRNA (serum)	Whole blood	10 miRNAs postulated to be involved in the regulation of genes associated with the neurometabolic cascade (miR-151-5p, miR-20a, miR-362-3p, miR-486, miR-505, miR-499, miR-625, miR-638, miR-381, miR-142-3p, miR-30d, miR-328, miR-27b, miR-92) were up-regulated in concussed individuals and progressive reduction at 24 h and 48 h post-concussion was observed.
Davies et al. (Davies et al., 2019)	41 athletes with a history of concussion, 15	45 males,	Not reported	25.1	Rugby Football Union HIA and SCAT-5.	miRNA (serum)	Whole blood	miR-502 implicated in the modulation of pathology in extra cranial diseases was

(continued on next page)

Table 3 (continued)

Study	Participants N	Sex	Ethnicity	Age (mean) [Years]	Concussion Diagnosis	Epigenetic mechanism	Tissue type	Main Findings
	controls without history of concussion.	10 females						downregulated for 7 days post-concussion compared to controls.
Papa et al. (Papa et al., 2019)	23 collegiate American Football athletes (9 athletes had a history of concussion and 2 in season concussions reported), 30 non-athlete controls with no history of concussion.	38 males, 15 females	Not reported	25.5	SAC-C (samples were collected pre- and post-season).	miRNA (serum)	Whole blood	Post-season 6 miRNAs (miR-195, miR-92a, miR-30d, miR-505*, miR-151-5p, and miR-362-3p) were elevated in the 2 athletes that sustained concussions over the season, only miRNA-92a was significantly higher post-season compared to non-concussed athletes. Higher levels of 4 miRNAs (miR-20a, miR-505*, miR-195, and miR-151-5p) were associated with lower neurocognitive scores in athletes compared to controls post-season.
Svingos et al. (Svingos et al., 2019)	27 collegiate athletes with a history of concussion.	11 males, 16 females	Mixed ethnicities	18.8	SCAT-3, SAC-C, (samples were collected pre-season and as soon as possible post-concussion (mean time of 6.8 h)).	miRNA (serum)	Whole blood	3 miRNAs (miR153-3p, miR223-3p, miR-let-7a-5p) associated with central nervous system dysfunction were upregulated post-concussion in male and female athletes.
Di Pietro et al. (Di Pietro et al., 2018)	28 rugby athletes with history of concussion, 16 rugby athlete controls with no history of concussion	44 males	Not reported	27.3	SCAT-3 (samples from concussed players were collected 48, 72 and > 120 h post-concussion).	miRNA (salivary)	Saliva	5 miRNAs (miR-27b-3p, let-7i-5p, miR-142-3p, miR-107, miR-135b-5p) were significantly upregulated in concussed athletes.
Johnson et al. (Johnson et al., 2018)	22 youths with acute concussion symptoms, 30 youth participants with prolonged concussion symptoms.	30 males, 22 females.	Mixed ethnicities	14.0	GCS and SCAT-3 within 14 days of injury.	miRNA (salivary)	Saliva	Levels of miRNAs (miR-320c-1, miR-133a-5p, miR-769-5p, let-7a-3p, and miR-1307-3p) associated with neuronal regulatory pathways were able to identify individuals with prolonged concussion symptoms compared to those with acute symptoms. Additionally, 3 miRNAs were associated with specific symptoms 4 weeks post-concussion; miR-320c-1 was associated with memory difficulty, miR-629 was associated with headaches, and let-7b-5p was associated with fatigue.
Pasinetti et al. (Pasinetti et al., 2012)	9 military veterans with history of concussion, 9 military control veterans without history of concussion.	13 males, 5 females	Mixed ethnicities	30.7	VAT-BIST (mean time between last deployment and recruitment to the study was 3.5 years).	small nucleolar RNA and miRNA (peripheral blood mononuclear cell isolation)	Whole blood	12 small nucleolar RNA (ACA48, ENSG199411, HBII-239, HBII-289, U15B, U27, U35A, U55, U56, U58B, U83A, U91) and 1 miRNA (hsa-miR-671-5p) were downregulated in veterans with a history of concussion compared to veterans with no history of concussion. 3 small nucleolar RNAs (HBII-289, ENSG199411 and U35A), could distinguish concussion cases in veteran participants.
Hicks et al. (Hicks et al., 2020)	251 participants with diagnosed concussion, 287	331 males,	Mixed ethnicities	18.3	Clinical assessments carried out < 14 days post-injury using the	ncRNA (salivary)	Saliva	A four-miRNA model (miR-34a-5p, miR-192-5p, miR-27a-5p, miR-4510) could

(continued on next page)

Table 3 (continued)

Study	Participants N	Sex	Ethnicity	Age (mean) [Years]	Concussion Diagnosis	Epigenetic mechanism	Tissue type	Main Findings
	controls with no recent history of concussion.	207 females			Zurich Concussion in Sport Group guidelines (samples collected < 3 days, 4–7 days, 8–14 days, 15–30 days and 31–60 days post-concussion).			differentiate concussed participants from non-concussed controls.
Miller et al. (Miller et al., 2022)	60 participants with diagnosed concussion.	32 males, 28 females	Mixed ethnicities	14.3	Clinical assessments carried out < 14 days post-injury using the Zurich Concussion in Sport Group guidelines, SAC-C and daily SCAT-5 (samples collected < 7 days, 7–14 days and 28 days post-concussion).	miRNA (salivary)	Saliva	13 miRNA levels (hsa-miR-95-3p, hsa-miR-301a-5p, hsa-miR-626, hsa-miR-548y, hsa-miR-203a-5p, hsa-miR-548e-5p, hsa-miR-585-3p, hsa-miR-378 h, hsa-miR-1323, hsa-miR-183-5p, hsa-miR-200a-3p, hsa-miR-888-5p, hsa-miR-199a-3p, hsa-miR-199b-3p) at 3 timepoints post-concussion were higher in concussed children with persisting symptoms after concussion compared to concussed children without persisting symptoms after concussion. At 0 days miR-32-5p was higher in concussed participants compared to controls. In concussed participants reporting higher symptom severity scores, at 7 days miR-32-5p was lower and at 28 days miR-142-3p and miR-223-3p were higher, compared to concussed participants who reported lower symptom severity scores. miR423-3p was higher among concussed participants < 6 hrs post-injury compared to controls.
Mitra et al. (Mitra et al., 2022)	28 participants with diagnosed concussion, 30 controls with no history of concussion.	28 males, 30 females	Not reported	31.0	GCS (samples were collected at 0 days, 7 days and 28 days post-concussion).	miRNA (plasma)	Whole blood	miR191 levels were elevated in participants with concussion compared to controls.
Mitra et al. (Mitra et al., 2023)	75 participants with diagnosed concussion, 44 controls with no history of concussion.	65 males, 54 females	Not reported	31.5	GCS (samples were collected at < 6 hrs of injury).	miRNA (plasma)	Whole blood	miR191 levels were elevated in participants with concussion compared to controls.
Tas et al. (Tas et al., 2020)	79 participants with diagnosed concussion, 92 controls with no history of acute trauma.	134 males, 37 females	Not reported	38.8	GCS (samples were collected < 24 hrs of admission).	miRNA (plasma)	Whole blood	From combining all timepoints, miRNA Let-7c-5p was upregulated in concussed athletes and 12 miRNAs were downregulated (miR181c-5p, miR-146a-5p, miR-200c-3p, miR-22-3p, miR-17-5p, miR26a-5p, miR-154-5p, miR-210-5p, miR-19b-3p, miR-16-5p, miR29a-3p, and miR-181c-3p).
Wyczechowska et al. (Wyczechowska et al., 2023)	57 collegiate American Football athletes (10 athletes sustained concussions and 11 non-concussed control athletes)	57 males	Mixed ethnicities	20.0	SCAT-3 and clinical assessments (samples were collected 1–2 hr and 18 hrs post-injury and end of season).	miRNA (serum)	Whole blood	Global methylation did not show significant concussion group versus control group differences. However, DNA methylation factors under the major depression pathway contributed to the prediction of the quality of life but not persistent post-concussive symptoms.
Duan et al. (Duan et al., 2021)	110 paediatric patients with a history of concussion, 87 controls with no history of concussion.	108 males, 89 females	Mixed ethnicities	14.9	Clinical assessments carried out 1–10 days post-injury using ACRM and the Zurich Concussion in Sport Group guidelines (samples collected 7.26 days post-concussion).	DNA methylation (850,000 CpG loci)	Saliva	

(continued on next page)

Table 3 (continued)

Study	Participants N	Sex	Ethnicity	Age (mean) [Years]	Concussion Diagnosis	Epigenetic mechanism	Tissue type	Main Findings
Bahado-Singh et al. (Bahado-Singh et al., 2020)	18 youths with a history of concussion, 18 youth controls without a history of concussion.	28 males, 8 females	White	12.8	GCS and SAC-C (8-hour time lapse after concussion and sample collection).	DNA methylation (850,000 CpG loci)	Whole blood	449 CpG sites were significantly differentially methylated (some methylation changes $\geq 10\%$) in concussed youths compared to youth controls. These CpG sites have been associated with; impaired brain function, cognition, memory, neurotransmission, intellectual disability, and behavioural change disorders.
Lee et al. (Lee et al., 2020)	11 collegiate students with a history of concussion, 14 collegiate student controls without history of concussion.	13 males, 12 females	Mixed ethnicities	28.7	Self-reported concussions (mean time lapse since the last concussion and sample collection was 7 years).	Global DNA methylation (5-mC%)	Whole blood	Higher blood global methylation ratio (5-mC%) in concussion cases (4.42 ± 0.43) than in controls (3.93 ± 0.03).
Schwab et al. (Schwab et al., 2019)	38 donated professional athlete brains with a history of concussion, 10 controls (5 healthy controls with no history of concussion, 5 AD cases with no history of concussion).	48 males	Not reported	54.6	Donated brains of professional contact sport athletes (>5 years exposure to repeated concussions).	Histone modification	Human brain tissue	Individuals with a history of concussion had a decreased level of H3K27Me3 in the context of DNA damage. This marker is considered marker of cellular senescence conferring susceptibility to brain dysfunction post-concussion.

OSU mTBI-ID, Ohio State university mTBI identification method; VCU rCDI, Virginia Commonwealth University retrospective concussion diagnosis interview; HIA, Head Injury Assessment; ACRM, American Congress of Rehabilitation Medicine; SCAT-3(5), Sport Concussion Assessment Tool-3(5); SAC-C, Standardised Assessment of Concussion score; GCS, Glasgow Coma Scale; CSF, Cerebral spinal fluid; AD, Alzheimer's Disease; PTSD, Post Traumatic Stress Disorder; VAT-BIST, Veteran traumatic brain injury screening tool, AFL, Australian Football League.

4.1. *sncRNAs*

The most extensively examined epigenetic modifiers in the studies from this review were miRNAs. miRNAs can act as post-transcriptional regulators of pathophysiological and functional cellular processes within the brain (Coolen and Bally-Cuif, 2009; Bhalala et al., 2013). Understanding the roles and functions of miRNA are rapidly expanding. However, for many miRNAs their full role and functions remain unclear. miRNAs are well suited to provide insight to the reactions of the injured brain following concussion as they can cross the blood-brain barrier and are stable in peripheral biofluids (Chen et al., 2016; García-Romero et al., 2017). In contrast, blood brain barrier disruption may be required for protein biomarkers such as, glial fibrillary acidic protein, neurofilament light chain protein and tau to migrate into peripheral tissues (Middeldorp and Hol, 2011) and can be degraded by endogenous proteases (Bhomia et al., 2016). Moderate and severe TBIs result in cell injury, and consequently protein biomarkers can be detected in peripheral tissues which can reliably be used to differentiate the severity of TBI (Honda et al., 2010). However, for the 'mild' form of TBI (concussion) cell injury may be subtle (Gonzalez et al., 2021). Thus, miRNAs and other sncRNAs may be more sensitive for diagnostic and prognostic purposes when compared to protein biomarkers, as their dysregulated expression in live cells may be more readily detected (Goldie et al., 2014).

4.1.1. Potential diagnostic miRNAs markers of concussion

It is possible that miRNAs could provide diagnostic insight to the management of concussion (Fig. 2). For example, the let-7 family and its

target genes play a role in neuroinflammation modulation (Lv et al., 2018). Di Pietro et al. reported that the upregulated let-7f-5p was able to provide good discrimination (AUC 0.80–0.89) between elite male rugby athletes with a history of concussion (36–68 h post-concussion) compared to those without a history of concussion (Di Pietro et al., 2021). Similarly, Svingos et al. observed an acute (~7 h post-concussion) upregulation of let-7f-5p in male and female collegiate athletes (Svingos et al., 2019).

The complex polygenic pathophysiology of concussion reflects the involvement of multiple miRNAs (Buckland et al., 2022;13:1107.; Barter and Foster, 2018). Therefore, the combination of multiple candidate miRNA biomarkers could provide additional objective diagnostic and prognostic insight to an individual's concussion status, alongside validated assessment tools. Di Pietro et al. demonstrated a 16 saliva sncRNA signature (let-7a-5p, miR-143-3p, miR-103a-3p, miR-34b-3p, RNU6-7, RNU6-45, snora57, snoU13.120, tRNA18Arg-CCT, U6-168, U6-428, U6-1249, Uco22c1g1, YRNA-255) could accurately identify concussed male elite rugby athletes from non-concussed participants immediately post-match (AUC = 0.91) and 36–48 h post-match (AUC = 0.94) (Di Pietro et al., 2021). Similarly, Hicks et al. demonstrated a single saliva miRNA ratio of miR-27a-5p/miR-30a-3p (logit (P) = 0.155 + 0.169) was able to provide a high accuracy (AUC = 0.810, sensitivity = 82.4 % specificity = 73.3 %) for identification of concussion status (Hicks et al., 2021). We propose that a subset of markers identified from this review may provide additional utility as a diagnostic signature although this requires further exploration.

Table 4
Differentially expressed miRNAs identified in multiple studies.

miRNA	Devoto et al. (2022)	Shultz et al. (2022)	Di Pietro et al. (Di Pietro V, O'Halloran P, Watson CN, Begum G, Acharjee A, Yakoub KM, et al. Unique diagnostic signatures of concussion in the saliva of male athletes: the Study of Concussion in Rugby Union through MicroRNAs (SCRUM). Br J Sports Med [Internet]. Br J Sports Med; 2021)	Fedorchak et al. (Fedorchak et al., 2021) [§]	Guedes et al. (Guedes et al., 2021)	Hicks et al. (Hicks et al., 2021)	Lusardi et al. (Lusardi et al., 2021)	Papa et al. (Papa et al., 2021)	Vorn et al. (Vorn et al., 2021)	Ghai et al. (Ghai et al., 2020)	Hicks et al. (Hicks et al., 2020)	Polito et al. (Polito et al., 2020)	Papa et al. (Papa et al., 2019)	Svingos et al. (Svingos et al., 2019)	Di Pietro et al. (Di Pietro et al., 2018)	Johnson et al. (Johnson et al., 2018)	Hicks et al. (Hicks et al., 2020)	Miller et al. (Miller et al., 2022) [§]	Mitra et al. (Mitra et al., 2022)	
Let 7a-5p			↑											↑						
miR-107			↑						↓							↑				
miR-1246*			↓		↓															
miR-125a-5p*							↓		↓											
miR-126-3p*									↓		↓									
miR-1307-3p																				
miR-135b-5p*			↑																	
miR-139-5p*	↓				↓															
miR-140-5p*							↓													
miR-142-3p									↓		↑		↑							↑
miR-148a-5p*																				
miR-148a-3p			↑																	
miR-151a-3p*									↓											
miR-181a-5p*																				
miR-182-5p																				
miR-18a-5p*	↑																			
miR-191-5p*							↓													
miR-192-5p																				
miR-199a-3p*																				
miR-204-5p*	↑				↑															
miR-20a-5p							↓													
miR-20b-5p							↓													
miR-21-5p			↑																	
miR-222-3p*																				
miR-223-3p																				
miR-27a-3p*		↓																		
miR-27b-3p																				
miR-28-3p*																				
miR-30d																				
miR-30e-5p*																				
miR-3190-3p*	↑				↑															
miR-362-3p							Mixed [#]	↑												
miR-372-3p*	↑				↑															
miR-423-5p																				
miR-505*																				
miR-615-5p*	↑				↑															
miR-92a																				

* miRNA differentially expressed in a consistent direction in at least two studies.

[#] Lusardi et al. (Lusardi et al., 2021) miR-362-3p was found in more military concussion participants than community controls but at lower levels. [§]Fedorchak et al. (Fedorchak et al., 2021), all participants had a history of concussion, the miRNA differential expression is between participants showing signs of persisting symptoms after concussion and those without symptoms. Papa et al. (Papa et al., 2021); Davies et al. (Davies et al., 2019); Pasinetti et al. (Pasinetti et al., 2012) were not included in the table as their identified miRNAs were not found in an additional study.




Fig. 2. Potential diagnostic and prognostic miRNAs markers of concussion.

4.1.2. Potential prognostic miRNAs markers of concussion

In addition, it is possible that miRNAs could provide some sort of prognostic use in the management of concussion (Fig. 2). For example, increased levels of miR-1307-3p could identify youths with persisting

symptoms post-concussion (4–8 weeks) (Johnson et al., 2018). Similarly, increased levels of miR-1307-3p were observed in adult military veterans 2–8 years after sustaining a concussion (Ghai et al., 2020). In recovery from concussion, Fedorchak et al. developed a signature using




Fig. 3. Ingenuity pathway analysis network of 10 miRNAs previously associated with concussion. Red-labelled miRNAs indicate increased expression post-concussion. Green-labelled miRNAs indicate decreased expression post-concussion. White-labelled miRNAs indicate inconclusive evidence to suggest direction of expression change post-concussion.

16 saliva sncRNAs (7 miRNAs, 1 snoRNA and 8 piRNA clusters) combined with age that could better predict persisting symptoms after concussion in youths (AUC = 0.83) than cognitive and balance testing (AUC = 0.79) (Fedorchak et al., 2021). However, combining sncRNAs with cognitive and balance testing yielded the greatest predictive accuracy (AUC = 0.86). Similarly, Johnson et al. developed a saliva miRNA model (miR-320c-1, miR-133a-5p, miR-769-5p, let-7a-3p, and miR-1307-3p) that could accurately identify youths with prolonged concussion symptoms (AUC = 0.86) (Johnson et al., 2018). Findings indicate that some miRNAs potentially have both diagnostic and prognostic properties (Fig. 2). Whilst these findings are still in the preliminary stages, individual biomarkers or combined models of epigenetic biomarkers may yet provide a useful corroborative diagnostic and prognostic tool for concussion management.

Ingenuity pathway analysis provides a valuable tool to help interpret and understand the significance of miRNA changes, and can identify gene signatures, which may provide putative biomarkers associated with concussion. With improved understanding of these pathway perturbations, clinicians may be better prepared to innovate targeted therapeutic interventions to improve patient outcomes post-concussion. Our work shows that a subset of 37 miRNAs have been identified from multiple studies, of which 10 were identified as having network interactions with 15 neuropathology associated genes. Further replication studies are needed to provide strong supporting evidence to confirm whether this gene signature is associated with concussion and further sncRNAs may be used to identify additional key networks, causal

relationships, and novel regulatory pathways that assist with the discovery of potential concussion biomarkers for diagnostic and prognostic purposes.

4.2. DNA methylation

Altered levels of global DNA methylation (average methylation status that occurs across the genome) have been associated with neuroinflammation and neurodegeneration such as, short-term memory loss (Tsankova et al., 2007; Tooby et al., 2023). Lee et al. demonstrated that concussion had a long-term effect on the global methylation ratio (5-mC %), as college athletes with a history of concussion (seven years post-concussion) had a higher blood global methylation ratio (4.42 ± 0.43) than controls with no history of concussion (3.93 ± 0.03) (Lee et al., 2020). In contrast, Duan et al. did not observe significant differences in global methylation between the youth concussion group compared to the control group (Duan et al., 2021). Differences in methods employed may account for these conflicting findings. Duan et al. collected saliva samples at 7.26 days post-concussion which, may not have been sufficient time for global methylation changes to occur in youths (Duan et al., 2021). The methylation profile of saliva could be different to that of blood, as it is more distant from the injury site. In addition, under/over estimation of global methylation may occur between studies as different assays were used (Kurdyukov and Methylation, 2016). Indeed, Duan et al. used Infinium Methylation EPIC array (Duan et al., 2021), whilst Lee et al. used enzyme-link immunosorbent assay (Lee et al., 2020).

However, global methylation ratios could be a useful tool in the future as an epigenetic marker of concussion recovery and potential associated neurodegeneration as changes in epigenetic marks have long-lasting effects within neurons (Tsankova, 2007). In addition, using artificial intelligence techniques such as deep learning, Bahado-Singh et al. demonstrated that 119 CpG methylation markers provided a classifier with good diagnostic accuracy ($AUC \geq 0.80$ – 0.89) for the detection of paediatric concussion (Bahado-Singh et al., 2020). In addition, combining epigenetic data with clinical data relating to concussion such as Sport Concussion Assessment Tool (SCAT) scores further increased the diagnostic accuracy of the concussion prediction model. Findings suggest that DNA methylation changes can be detected years post-concussion. Thus, providing long-term markers of concussion that could be used to monitor recovery may present target sites for therapeutic development and stratified intervention.

4.3. Histone modification

There is a paucity of data about the effects of concussion on histone modifications and the potential influences on outcomes and recovery post-concussion. Findings from Schwab et al. suggest that an inability to effectively repair DNA could impact the long-term effects of concussion (Schwab et al., 2019). Histone H3K27Me3 modifications are reduced in the context of DNA damage in individuals with a history of concussion (Schwab et al., 2019). Reduced expression of this histone mark in glial cells suggests activation of senescence through activation of p16 and p21 and is linked with neurodegenerative and cognitive decline (Schwab et al., 2019). Reduced expression of H3K27Me3 may also confer altered DNA damage responses potentially influencing recovery and risk of neurodegenerative disease post-concussion. It is plausible that a combination of histone modification, DNA methylation, and sncRNA expression contribute to neurodegenerative pathology and symptomology associated with concussion.

4.4. Clinical applications

In-depth neurological examination and sophisticated neuroimaging are not widely used to diagnose concussion, especially in youth and sub-elite level sports. Typically, clinicians rely upon subjective symptom reports to inform diagnosis, which could contribute to the under-diagnosis of concussions (Broglia et al., 2017; Meehan et al., 2013; McCrea et al., 2004). Individual epigenetic profiles in response to concussion could have the capacity to provide additional objective information to clinicians when diagnosing, managing recovery, and informing return to play processes in conjunction with current methods. Such profiles could be correlated against data derived from the developing use of impact markers, such as sensors placed in mouthguards (Tooby et al., 2023), to inform what level of force correlates with significant change in neurological cellular function – both within the general athletic population and individuals.

Epigenetic markers could be used to confirm that an individual has not sustained a concussion, thus providing reassurance of a negative concussion test. Another important clinical application is that miRNAs could serve as potential biomarkers to aid prognosis of concussion to identify individuals more at risk for persistent post-concussive symptoms and those who will have a typical recovery. Individualised therapy could be honed with the additional information derived from epigenetic markers, for example, early detection of markers associated with post-concussion chronic migraines could inform triptan therapy (Capi et al., 2020). Aerobic exercise reduces elevated post-TBI levels of miR-21 and facilitates cognitive improvements post-TBI in mice (Hu et al., 2015). Such findings suggest that monitoring of miRNAs as part of return to play protocols may inform the knowledge of likely recovery times for an individual. Sub-concussive head injuries are a growing concern due to the associations with neurodegenerative conditions such as chronic traumatic encephalopathy [92–93]. Future research

investigating epigenetic markers sensitive enough to detect sub-concussive insults to the brain could provide a useful tool to inform protective protocols for individuals. In addition, markers of sub-concussion could provide insight into the long-term effects on brain health. However, these clinical applications are currently speculative, as findings highlighted in this review need to be sufficiently validated before future guidance for clinical care and practice.

4.5. Limitations

This review is limited by challenges in combining and comparing methods utilised such as sample tissue types, populations, mechanisms of concussion, injury diagnoses, sample collection times. The assays platforms used to measure epigenetic alterations were also non-uniform. In addition, we recognise that epigenetic modifications exhibit population differences (Barfield et al., 2014) therefore, the use of heterogeneous cohorts in studies could affect findings and may not replicate across different populations. For miRNAs that are upregulated shortly after concussion and downregulated years later, future studies could investigate whether there is a long-term compensation mechanism which occurs and remains even after reaching pre-injury baseline levels. Longitudinal studies that monitor the dynamic sncRNA responses post-concussion could be used to provide both short and long-term prognostic insight into recovery and potential neuropathological consequences of concussion. Concussion-related epigenetic responses are complex and can be influenced by many additional confounding factors outside of the injury mechanism including demographics, ethnicity, age, sex, lifestyle, comorbidities, and medications (Alegria-Torres et al., 2011). Controlling for confounding factors in relation to concussion injuries will provide further insight into the molecular mechanisms of concussion processes and improve the management of this injury.

5. Conclusion

The emerging evidence highlights complex epigenetic variation that influences concussion pathophysiology. This knowledge is contributing to our understanding of genes, pathways and biological networks involved in response to concussion. Indeed, supplementary epigenetic information has the potential to assist in identification of concussions and to assist in individualised concussion management strategies. In our opinion, salivary candidate miRNAs could prove the most practical and insightful tool in managing an individual's concussion status. As indicated by our IPA, it is likely that combining data from multiple concussion-associated miRNAs reflects the complexity of this injury and could increase the accuracy of a practical tool for concussion screening and management. However, the research is still in its infancy and significant knowledge gaps exist. Each concussion is a unique injury and is influenced by numerous genetic and environmental factors. This creates many challenges to consider when investigating epigenetic markers for future studies. However, for epigenetic markers of concussion to have a practical application, they will need to demonstrate replicability across studies with consistent findings evidenced.

CRediT authorship contribution statement

Mark R. Antrobus: Writing – original draft. **Terun Desai:** Writing – original draft. **David Young:** Writing – original draft. **Lee Machado:** Writing – original draft. **William J. Ribbans:** Writing – original draft. **Louis Y. El Khoury:** Writing – original draft. **Jon Brazier:** Writing – original draft.

Funding

This research received no external funding.

All authors contributed to the initial development of the review and search criteria and collectively interpreted the results of the systematic

review and meta-analysis. MRA, TD and JB contributed to the implementation of the search strategy and application of the inclusion/exclusion criteria and quality scoring system. All authors drafted the manuscript and all authors contributed to editing and revising the manuscript and approved the final version prior to submission.

Ethics Approval: Not applicable.

Consent to Participate: Not applicable.

Consent for Publication: Not applicable.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data will be made available on request.

References

- Abrahams, S., McFie, S., Patricios, J., Posthumus, M., September, A.V., 2014. Risk factors for sports concussion: an evidence-based systematic review. Available from Br J Sports Med [Internet]. BMJ Publishing Group Ltd and British Association of Sport and Exercise Medicine. 48, 91–97.
- Alegria-Torres, J.A., Baccarelli, A., Bollati, V., 2011. Epigenetics and lifestyle. *Epigenomics* [Internet]. NIH Public Access. 3, 267–277 [cited 2023 Jan 17]. Available from: [/pmc/articles/PMC3752894/](https://pubmed.ncbi.nlm.nih.gov/31628932/).
- Antrobus, M.R., Brazier, J., Stebbings, G.K., Day, S.H., Heffernan, S.M., Kilduff, L.P., et al., 2021. Genetic factors that could affect concussion risk in elite rugby [cited 2021 Feb 18];9:19. Available from: Sports [Internet]. MDPI AG 9, 19 <https://www.mdpi.com/2075-4663/9/2/19>.
- Bahado-Singh, R.O., Vishweswaraiah, S., Er, A., Aydas, B., Turkoglu, O., Taskin, B.D., et al., 2020. Artificial Intelligence and the detection of pediatric concussion using epigenomic analysis. *Brain Res* [Internet] [cited 2023 Jan 16];1726. Available from: *Brain Res* <https://pubmed.ncbi.nlm.nih.gov/31628932/>.
- Bannister AJ, Kouzarides T. Regulation of chromatin by histone modifications. *Cell Res* [Internet]. Nature Publishing Group; 2011 [cited 2023 Jan 13];21:381–95. Available from: <https://www.nature.com/articles/cr201122>.
- Barfield, R.T., Almlí, L.M., Kilaru, V., Smith, A.K., Mercer, K.B., Duncan, R., et al., 2014. Accounting for population stratification in DNA methylation studies [cited 2023 Jan 17]. Available from *Genet Epidemiol* [Internet] *Genet Epidemiol*. 38, 231–241.
- Barkhoudarian, G., Hovda, D.A., Giza, C.C., 2016. The molecular pathophysiology of concussive brain injury – an update [cited 2019 Oct 16]; Available from: *Phys Med Rehabil Clin N Am* [Internet]. 27, 373–393 <http://www.ncbi.nlm.nih.gov/pubmed/27154851>.
- Barter JD, Foster TC. Aging in the Brain: New Roles of Epigenetics in Cognitive Decline. <https://doi.org/10.1177/1073858418780971> [Internet]. SAGE PublicationsSage CA: Los Angeles, CA; 2018 [cited 2023 Jan 13];24:516–25. Available from: <https://journals.sagepub.com/doi/10.1177/1073858418780971>.
- Bhalala OG, Srikanth M, Kessler JA. The emerging roles of microRNAs in CNS injuries. *Nat Rev Neurol* [Internet]. Nat Rev Neurol; 2013 [cited 2023 Jan 19];9:328–39. Available from: <https://pubmed.ncbi.nlm.nih.gov/23588363/>.
- Bhomia M, Balakathiresan NS, Wang KK, Papa L, Maheshwari RK. A Panel of Serum miRNA Biomarkers for the Diagnosis of Severe to Mild Traumatic Brain Injury in Humans. *Sci Reports* 2016 61 [Internet]. Nature Publishing Group; 2016 [cited 2023 Jan 30];6:1–12. Available from: <https://www.nature.com/articles/srep28148>.
- Bieniek, K.F., Blessing, M.M., Heckman, M.G., Diehl, N.N., Serie, A.M., Paolini, M.A., et al., 2020. Association between contact sports participation and chronic traumatic encephalopathy: a retrospective cohort study [cited 2022 Oct 4];30:63–74. Available from: *Brain Pathol* [Internet] *Brain Pathol* <https://pubmed.ncbi.nlm.nih.gov/31199537/>.
- Broglio, S.P., McCrea, M., McAllister, T., Harezlak, J., Katz, B., Hack, D., et al., 2017. A national study on the effects of concussion in collegiate athletes and US military service academy members: The NCAA-DoD concussion assessment, research and education (CARE) consortium structure and methods. *sports med* [Internet] [cited 2022 Oct 3] Available from *Sports Med* 47, 1437–1451.
- Buckland, M.E., Affleck, A.J., Pearce, A.J., Suter, C.M., 2022. Chronic traumatic encephalopathy as a preventable environmental disease. *Front Neurol*. *Frontiers Media* s.a. 13, 1107.
- Capi, M., Pomes, L.M., Andolina, G., Curto, M., Martelletti, P., Lionetto, L., 2020. Persistent post-traumatic headache and migraine: pre-clinical comparisons. *Int J Environ Res Public Health* [Internet] Multidisciplinary Digital Publishing Institute (MDPI). 17 [cited 2023 Mar 16]; Available from: [/pmc/articles/PMC7177371/](https://pubmed.ncbi.nlm.nih.gov/3177371/).
- Chen CC, Liu L, Ma F, Wong CW, Guo XE, Chacko J V., et al. Elucidation of Exosome Migration across the Blood-Brain Barrier Model In Vitro. *Cell Mol Bioeng* [Internet]. Cell Mol Bioeng; 2016 [cited 2023 Jan 18];9:509–29. Available from: <https://pubmed.ncbi.nlm.nih.gov/28392840/>.
- Coolen M, Bally-Cuif L. MicroRNAs in brain development and physiology. *Curr Opin Neurobiol* [Internet]. *Curr Opin Neurobiol*; 2009 [cited 2023 Jan 19];19:461–70. Available from: <https://pubmed.ncbi.nlm.nih.gov/19846291/>.
- Dana H, Mahmoodi Chalbatani G, Mahmoodzadeh H, Karimloo R, Rezaiean O, Moradzadeh A, et al. Molecular Mechanisms and Biological Functions of siRNA. *Int J Biomed Sci* [Internet]. Master Publishing Group; 2017 [cited 2024 Apr 16];13:57. Available from: [/pmc/articles/PMC5542916/](https://pubmed.ncbi.nlm.nih.gov/35542916/).
- Davies D, Yakoub KM, Scarpa U, Bentley C, Grey M, Hammond D, et al. Serum miR-502: A potential biomarker in the diagnosis of concussion in a pilot study of patients with normal structural brain imaging. *J Concussion* [Internet]. SAGE PublicationsSage UK: London, England; 2019 [cited 2023 Feb 10];3:1–13. Available from: <https://journals.sagepub.com/doi/full/10.1177/2059700219886190>.
- Devoto C, Guedes VA, Lai C, Leete JJ, Mithani S, Edwards K, et al. Remote blast-related mild traumatic brain injury is associated with differential expression of exosomal microRNAs identified in neurodegenerative and immunological processes. *Brain Inj* [Internet]. *Brain Inj*; 2022 [cited 2023 Jan 27];36:652–61. Available from: <https://pubmed.ncbi.nlm.nih.gov/35322723/>.
- Dewan, M.C., Rattani, A., Gupta, S., Baticulon, R.E., Hung, Y.C., Punchak, M., et al., 2018. Estimating the global incidence of traumatic brain injury [cited 2022 Oct 3] Available from: *J Neurosurg* [Internet]. American Association of Neurological Surgeons 130, 1080–1097 <https://thejns.org/view/journals/j-neurosurg/130/4/article-p1080.xml>.
- Di Pietro V, O'Halloran P, Watson CN, Begum G, Acharjee A, Yakoub KM, et al. Unique diagnostic signatures of concussion in the saliva of male athletes: the Study of Concussion in Rugby Union through MicroRNAs (SCRUM). *Br J Sports Med* [Internet]. *Br J Sports Med*; 2021 [cited 2023 Jan 18];55:1395–404. Available from: <https://pubmed.ncbi.nlm.nih.gov/33757972/>.
- Di Pietro, V., Porto, E., Ragusa, M., Barbagallo, C., Davies, D., Forcione, M., et al., 2018. Salivary MICORNAs: diagnostic markers of mild traumatic brain injury in contact-sport. *Front Mol Neurosci* [Internet] *Frontiers Media* s.a. [cited 2023 Jan 24];11: 336. Available from: [/pmc/articles/PMC6109773/](https://pubmed.ncbi.nlm.nih.gov/31628932/).
- Duan K, Mayer AR, Shaff NA, Chen J, Lin D, Calhoun VD, et al. DNA methylation under the major depression pathway predicts pediatric quality of life four-month post-pediatric mild traumatic brain injury. *Clin Epigenetics* [Internet]. BioMed Central Ltd; 2021 [cited 2023 Jan 13];13:1–17. Available from: <https://clinicalepigeneticsjournal.biomedcentral.com/articles/10.1186/s13148-021-01128-z>.
- Fedorchak G, Rangnekar A, Onks C, Loeffert AC, Loeffert J, Olympia RP, et al. Saliva RNA biomarkers predict concussion duration and detect symptom recovery: a comparison with balance and cognitive testing. *J Neurol* [Internet]. *J Neurol*; 2021 [cited 2023 Jan 27];268:4349–61. Available from: <https://pubmed.ncbi.nlm.nih.gov/34028616/>.
- García-Romero N, Carrión-Navarro J, Esteban-Rubio S, Lázaro-Ibáñez E, Peris-Celda M, Alonso MM, et al. DNA sequences within glioma-derived extracellular vesicles can cross the intact blood-brain barrier and be detected in peripheral blood of patients. *Oncotarget* [Internet]. *Oncotarget*; 2017 [cited 2023 Jan 18];8:1416–28. Available from: <https://pubmed.ncbi.nlm.nih.gov/27902458/>.
- Ghai V, Fallen S, Baxter D, Scherler K, Kim TK, Zhou Y, et al. Alterations in Plasma microRNA and Protein Levels in War Veterans with Chronic Mild Traumatic Brain Injury. *J Neurotrauma* [Internet]. Mary Ann Liebert, Inc.; 2020 [cited 2023 Jan 27];37:1430. Available from: [/pmc/articles/PMC7249467/](https://pubmed.ncbi.nlm.nih.gov/3249467/).
- Goldie BJ, Dun MD, Lin M, Smith ND, Verrills NM, Dayas C V., et al. Activity-associated miRNA are packaged in Map1b-enriched exosomes released from depolarized neurons. *Nucleic Acids Res* [Internet]. *Nucleic Acids Res*; 2014 [cited 2023 Mar 9];42:9195–208. Available from: <https://pubmed.ncbi.nlm.nih.gov/25053844/>.
- Gonzalez, A.C., Kim, M., Keser, Z., Ibrahim, L., Singh, S.K., Ahmad, M.J., et al., 2021. Diffusion tensor imaging correlates of concussion related cognitive impairment. *Front Neurol* [Internet] *Frontiers Media* s.a. [cited 2023 Mar 10];12: 639179. Available from: [/pmc/articles/PMC8180854/](https://pubmed.ncbi.nlm.nih.gov/34690777/).
- Guedes, V.A., Lai, C., Devoto, C., Edwards, K.A., Mithani, S., Sass, D., et al., 2021. Extracellular vesicle proteins and MicroRNAs are linked to chronic post-traumatic stress disorder symptoms in service members and veterans with mild traumatic brain injury [cited 2023 Jan 27];12. Available from: *Front Pharmacol* [Internet]. *Front Pharmacol* <https://pubmed.ncbi.nlm.nih.gov/34690777/>.
- Hicks SD, Olympia RP, Onks C, Kim RY, Zhen KJ, Fedorchak G, et al. Saliva microRNA Biomarkers of Cumulative Concussion. *Int J Mol Sci* [Internet]. Multidisciplinary Digital Publishing Institute (MDPI); 2020 [cited 2024 Jan 11];21:1–16. Available from: [/pmc/articles/PMC7589940/](https://pubmed.ncbi.nlm.nih.gov/34461327/).
- Hicks SD, Onks C, Kim RY, Zhen KJ, Loeffert J, Loeffert AC, et al. Refinement of saliva microRNA biomarkers for sports-related concussion. *J Sport Heal Sci* [Internet]. *J Sport Health Sci*; 2021 [cited 2023 Jan 27];21:S2095-2546. Available from: <https://pubmed.ncbi.nlm.nih.gov/34461327/>.
- Hicks, S.D., Onks, C., Kim, R.Y., Zhen, K.J., Loeffert, J., Loeffert, A.C., et al., 2020. Diagnosing mild traumatic brain injury using saliva RNA compared to cognitive and balance testing. *Clin Transl Med* [Internet]. Wiley-Blackwell [cited 2023 Jan 27];10: e197. Available from: [/pmc/articles/PMC7533415/](https://pubmed.ncbi.nlm.nih.gov/34461327/).
- Honda M, Tsuruta R, Kaneko T, Kasaoka S, Yagi T, Todani M, et al. Serum glial fibrillary acidic protein is a highly specific biomarker for traumatic brain injury in humans compared with S-100B and neuron-specific enolase. *J Trauma* [Internet]. *J Trauma*; 2010 [cited 2023 Mar 9];69:104–9. Available from: <https://pubmed.ncbi.nlm.nih.gov/20093985/>.
- Hu, T., Zhou, F.J., Chang, Y.F., Li, Y.S., Liu, G.C., Hong, Y., et al., 2015. miR21 is associated with the cognitive improvement following voluntary running wheel exercise in TBI mice [cited 2023 Mar 16] Available from *J Mol Neurosci* [Internet] *J Mol Neurosci*. 57, 114–122.

- Iverson, G.L., Gardner, A.J., Terry, D.P., Ponsford, J.L., Sills, A.K., Broshek, D.K., et al., 2017. Predictors of clinical recovery from concussion: a systematic review [cited 2022 Oct 3];51:941–8. Available from: [Br J Sports Med \[internet\] Br J Sports Med https://pubmed.ncbi.nlm.nih.gov/28566342/](https://pubmed.ncbi.nlm.nih.gov/28566342/).
- Johnson, J.J., Loeffert, A.C., Stokes, J., Olympia, R.P., Bramley, H., Hicks, S.D., 2018. Association of salivary MICRORNA changes with prolonged concussion symptoms. *JAMA Pediatr* [Internet]. American Medical Association [cited 2023 Jan 27];172: 73. Available from: [/pmc/articles/PMC5833519/](https://pubmed.ncbi.nlm.nih.gov/3017352/).
- June, D., Williams, O.A., Huang, C.W., An, Y., Landman, B.A., Davatzikos, C., et al., 2023. Lasting consequences of concussion on the aging brain: Findings from the Baltimore Longitudinal Study of Aging. *Neuroimage* [Internet]. NIH Public Access 221, 117182 [cited 2022 Oct 3]; Available from: [/pmc/articles/PMC7848820/](https://pubmed.ncbi.nlm.nih.gov/37488820/).
- Kim, S.K., Roche, M.D., Fredericson, M., Dragoo, J.L., Horton, B.H., Avins, A.L., et al., 2020. A genome-wide association study for concussion risk [cited 2021 Feb 22]; Available from: [Med Sci Sport Exerc \[internet\] Ovid Technologies \(wolters Kluwer Health\) https://pubmed.ncbi.nlm.nih.gov/33017352/](https://pubmed.ncbi.nlm.nih.gov/33017352/).
- Kurdyukov S, Bullock M. DNA Methylation Analysis: Choosing the Right Method. *Biology (Basel)* [Internet]. Multidisciplinary Digital Publishing Institute (MDPI); 2016 [cited 2023 Jan 13];5:1–21. Available from: [/pmc/articles/PMC4810160/](https://pubmed.ncbi.nlm.nih.gov/33017352/).
- Lee, H., Lee, S., Black, I., Salado, L., Estrada, J., Isla, K., 2020. Long-term impact of mild traumatic brain injuries on multiple functional outcomes and epigenetics: a pilot study with college students [cited 2023 Jan 13];10: 4131. Available from: [Appl Sci \[internet\]. Multidisciplinary Digital Publishing Institute https://www.mdpi.com/2076-3417/10/12/4131.htm](https://pubmed.ncbi.nlm.nih.gov/34580583/).
- Lefevre-Dognin, C., Cogné, M., Perdrieau, V., Granger, A., Heslot, C., Azouvi, P., 2021. Definition and epidemiology of mild traumatic brain injury. *Neurochirurgie. Elsevier Masson* 67, 218–221.
- Li, S., Lei, Z., Sun, T., 2022. The role of microRNAs in neurodegenerative diseases: a review [cited 2024 Apr 16];39:53–83. Available from: [Cell Biol Toxicol \[internet\]. Springer https://link.springer.com/article/10.1007/s10565-022-09761-x](https://link.springer.com/article/10.1007/s10565-022-09761-x).
- Lusardi, T.A., Sandau, U.S., Sakhanenko, N.A., Baker, S.C.B., Wiedrick, J.T., Lapidus, J. A., et al., 2021. Cerebrospinal fluid microRNA changes in cognitively normal veterans with a history of deployment-associated mild traumatic brain injury [cited 2023 Jan 27];15. Available from: [Front Neurosci \[internet\]. Front Neurosci https://pubmed.ncbi.nlm.nih.gov/34580583/](https://pubmed.ncbi.nlm.nih.gov/34580583/).
- Lust, C.A.C., Mountjoy, M., Robinson, L.E., Oliver, J.M., Ma, D.W.L., 2020. Sports-related concussions and subconcussive impacts in athletes: incidence, diagnosis, and the emerging role of EPA and DHA [cited 2022 Oct 4];45:886–92. Available from: [Appl Physiol Nutr Metab \[internet\] Appl Physiol Nutr Metab https://pubmed.ncbi.nlm.nih.gov/32119565/](https://pubmed.ncbi.nlm.nih.gov/32119565/).
- Lv J, Zeng Y, Qian Y, Dong J, Zhang Z, Zhang J. MicroRNA let-7c-5p improves neurological outcomes in a murine model of traumatic brain injury by suppressing neuroinflammation and regulating microglial activation. *Brain Res* [Internet]. *Brain Res*; 2018 [cited 2023 Jan 18];1685:91–104. Available from: <https://pubmed.ncbi.nlm.nih.gov/29408500/>.
- Maas, A.I.R., Menon, D.K., David Adelson, P.D., Andelic, N., Bell, M.J., Belli, A., et al., 2017. Traumatic brain injury: integrated approaches to improve prevention, clinical care, and research [cited 2022 Jun 27]; Available from: [Lancet Neurol \[internet\]. Elsevier 16, 987–1048 http://www.thelancet.com/article/S147444221730371X/fulltext](http://www.thelancet.com/article/S147444221730371X/fulltext).
- Maas, A.I.R., Menon, D.K., Manley, G.T., Abrams, M., Åkerlund, C., Andelic, N., et al., 2022. Traumatic brain injury: progress and challenges in prevention, clinical care, and research [cited 2022 Oct 3]; Available from: [Lancet Neurol \[internet\]. 1–19.](https://pubmed.ncbi.nlm.nih.gov/35712307/)
- McCrea, M., Hammeke, T., Olsen, G., Leo, P., Guskiewicz, K., 2004. Unreported concussion in high school football players: implications for prevention. *Clin J Sport Med. Lippincott Williams and Wilkins* 14, 13–17.
- McGowan J, Sampson M, Salzwedel M, Cogo E, Foerster V, Lefebvre C. CADTH Methods and Guidelines PRESS Peer Review of Electronic Search Strategies: 2015 Guideline Explanation and Elaboration (PRESS E&E). Ottawa; 2016.
- Mckee, A.C., Daneshvar, D.H., 2015. The neuropathology of traumatic brain injury [cited 2022 Oct 3];127:45–66. Available from: [Handb Clin Neurol \[internet\] Handb Clin Neurol https://pubmed.ncbi.nlm.nih.gov/25702209/](https://pubmed.ncbi.nlm.nih.gov/25702209/).
- Meehan, W.P., Mannix, R.C., Oebrien, M.J., Collins, M.W., 2013. The prevalence of undiagnosed concussions in athletes. *Clin J Sport Med* [internet]. NIH Public Access 23, 342 [cited 2022 Oct 3]; Available from: [/pmc/articles/PMC3758800/](https://pubmed.ncbi.nlm.nih.gov/21219963/).
- Middeldorp J, Hol EM. GFAP in health and disease. *Prog Neurobiol* [Internet]. *Prog Neurobiol*; 2011 [cited 2023 Mar 13];93:421–43. Available from: <https://pubmed.ncbi.nlm.nih.gov/21219963/>.
- Miller KE, MacDonald JP, Sullivan L, Venkata LPR, Shi J, Yeates KO, et al. Salivary miRNA Expression in Children With Persistent Post-concussive Symptoms. *Front Public Heal* [Internet]. *Front Public Health*; 2022 [cited 2024 Jan 11];10. Available from: <https://pubmed.ncbi.nlm.nih.gov/35712307/>.
- Mitra B, Reyes J, O'Brien WT, Surendran N, Carter A, Bain J, et al. Micro-RNA levels and symptom profile after mild traumatic brain injury: A longitudinal cohort study. *J Clin Neurosci* [Internet]. *J Clin Neurosci*; 2022 [cited 2024 Jan 11];95:81–7. Available from: <https://pubmed.ncbi.nlm.nih.gov/34929656/>.
- Mitra B, Major BP, Reyes J, Surendran N, Bain J, Giesler LP, et al. MicroRNA biomarkers for diagnosis of mild traumatic brain injury and prediction of persistent symptoms: A prospective cohort study. *J Clin Neurosci* [Internet]. *J Clin Neurosci*; 2023 [cited 2024 Jan 11];115:38–42. Available from: <https://pubmed.ncbi.nlm.nih.gov/37480731/>.
- Moore LD, Le T, Fan G. DNA Methylation and Its Basic Function. *Neuropsychopharmacol* 2013 381 [Internet]. Nature Publishing Group; 2012 [cited 2023 Jan 13];38:23–38. Available from: <https://www.nature.com/articles/npp2012112>.
- Nowinski, C.J., Bureau, S.C., Buckland, M.E., Curtis, M.A., Daneshvar, D.H., Faull, R.L. M., et al., 2022. Applying the Bradford Hill criteria for causation to repetitive head impacts and chronic traumatic encephalopathy [cited 2023 Mar 23];13. Available from: [Front Neurol \[internet\]. Front Neurol https://pubmed.ncbi.nlm.nih.gov/35937061/](https://pubmed.ncbi.nlm.nih.gov/35937061/).
- Page MJ, Moher D, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. *Br Med J* [Internet]. British Medical Journal Publishing Group; 2021 [cited 2023 Feb 7];372. Available from: <https://www.bmj.com/content/372/bmj.n160>.
- Panena WJ, Gardner AJ, Dretsch MN, Crynen GC, Crawford FC, Iverson GL. Systematic Review of Genetic Risk Factors for Sustaining a Mild Traumatic Brain Injury [Internet]. *J Neurotrauma. Mary Ann Liebert Inc.*; 2017 [cited 2020 Apr 7]. p. 2093–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28100103>.
- Papa L, Slobounov SM, Breiter HC, Walter A, Bream T, Seidenberg P, et al. Elevations in MicroRNA Biomarkers in Serum Are Associated with Measures of Concussion, Neurocognitive Function, and Subconcussive Trauma over a Single National Collegiate Athletic Association Division I Season in Collegiate Football Players. *J Neurotrauma* [Internet]. Mary Ann Liebert, Inc.; 2019 [cited 2023 Feb 10];36: 13Available from: [/pmc/articles/PMC6909756/](https://pubmed.ncbi.nlm.nih.gov/33017352/).
- Papa L, Johnson B, Walter AE, Wilkes JR, Knollmann-Ritschel B, Bhomia M, et al. Decreases in Dorsal Cervical Spinal Cord White Matter Tract Integrity Are Associated with Elevated Levels of Serum MicroRNA Biomarkers in NCAA Division I Collegiate Football Players. *Neurotrauma Reports* [Internet]. Mary Ann Liebert Inc; 2021 [cited 2023 Feb 10];2:476–87. Available from: <https://www.liebertpub.com/doi/10.1089/neur.2021.0036>.
- Pasinetti, G.M., Ho, L., Dooley, C., Abbi, B., Lange, G., 2012. Select non-coding RNA in blood components provide novel clinically accessible biological surrogates for improved identification of traumatic brain injury in OEF/OIF Veterans. *Am J Neurodegener Dis* [internet]. e-Century Publishing Corporation 1, 98. Available from: [/pmc/articles/PMC3560446/](https://pubmed.ncbi.nlm.nih.gov/3560446/).
- Polito, F., Famà, F., Oteri, R., Raffa, G., Vita, G., Conti, A., et al., 2020. Circulating miRNAs expression as potential biomarkers of mild traumatic brain injury [cited 2023 Jan 30] Available from [Mol Biol Rep \[internet\] Mol Biol Rep. 4747, 29412941–29492949.](https://pubmed.ncbi.nlm.nih.gov/32942949/)
- Ponsford, J., Willmott, C., Rothwell, A., Cameron, P., Kelly, A.M., Nelms, R., et al., 2000. Factors influencing outcome following mild traumatic brain injury in adults. Available from: [cited 2022 Oct 4] *J Int Neuropsychol Soc* [internet]. *J Int Neuropsychol Soc* 6, 568–579. <https://pubmed.ncbi.nlm.nih.gov/10932476/>.
- Raikes, A.C., Athey, A., Alfonso-Miller, P., Killgore, W.D.S., Grandner, M.A., 2019. Insomnia and daytime sleepiness: risk factors for sports-related concussion. *Sleep Med. Elsevier* b.v. 58, 66–74.
- Russell, E.R., Mackay, D.F., Lyall, D., Stewart, K., MacLean, J.A., Robson, J., et al., 2022. Neurodegenerative disease risk among former international rugby union players [cited 2022 Oct 5] Available from [J Neurol Neurosurg Psychiatry \[internet\]. BMJ Publishing Group Ltd 1–7.](https://pubmed.ncbi.nlm.nih.gov/3704304/)
- Sato, K., Takayama, K.I., Inoue, S., 2023. Role of piRNA biogenesis and its neuronal function in the development of neurodegenerative diseases. *Front Aging Neurosci* [Internet]. *Frontiers Media SA* [cited 2024 Apr 16];15. Available from: [/pmc/articles/PMC10191213/](https://pubmed.ncbi.nlm.nih.gov/410191213/).
- Schwab, N., Grenier, K., Hazrati, L.N., 2019. DNA repair deficiency and senescence in concussed professional athletes involved in contact sports. *Acta Neuropathol Commun* [internet] *BioMed Central* 7, 1–23 [cited 2023 Jan 12] Available from: [/pmc/articles/PMC6857343/](https://pubmed.ncbi.nlm.nih.gov/36857343/).
- Shultz, S.R., Taylor, C.J., Aggio-Bruce, R., O'Brien, W.T., Sun, M., Gioanca, A.V., et al., 2022. Decrease in plasma miR-27a and miR-221 after concussion in Australian football players. *Biomark Insights* [internet]. SAGE Publications [cited 2023 Jan 30]; 17. Available from: [/pmc/articles/PMC38891921/](https://pubmed.ncbi.nlm.nih.gov/38891921/).
- Stepanov, G.A., Filippova, J.A., Komissarov, A.B., Kuligina, E.V., Richter, V.A., Semenov, D.V., et al., 2015;2015. Regulatory role of Small nucleolar RNAs in human diseases. *Biomed Res Int. Hindawi Publishing Corporation*.
- Svingos AM, Asken BM, Bauer RM, DeKosky ST, Hromas GA, Jaffee MS, et al. Exploratory study of sport-related concussion effects on peripheral micro-RNA expression. *Brain Inj* [Internet]. *Brain Inj*; 2019 [cited 2023 Jan 18];33:1–7. Available from: <https://pubmed.ncbi.nlm.nih.gov/30704304/>.
- Tas, D., Kaplan, O., Sogut, O., 2020. Validity of serum miRNA 93 and miRNA 191 to reduce unnecessary computed tomography in patients with mild head trauma. *J Clin Med Res* [Internet]. *Elmer Press* 12, 579 [cited 2024 Jan 11] Available from: [/pmc/articles/PMC7430915/](https://pubmed.ncbi.nlm.nih.gov/37430915/).
- Tooby J, Woodward J, Tucker R, Jones B, Falvey É, Salmon D, et al. Instrumented Mouthguards in Elite-Level Men's and Women's Rugby Union: The Incidence and Propensity of Head Acceleration Events in Matches. *Sport Med* [Internet]. Springer Science and Business Media Deutschland GmbH; 2023 [cited 2024 Apr 4];1–12. Available from: <https://link.springer.com/article/10.1007/s40279-023-01953-7>.
- Tsankova, N., Renthal, W., Kumar, A., Nestler, E.J., 2007. Epigenetic regulation in psychiatric disorders [cited 2023 Mar 14] Available from [Nat Rev Neurosci \[internet\] Nature Publishing Group. 8, 355–367.](https://pubmed.ncbi.nlm.nih.gov/1735367/)
- Vorn R, Suarez M, White JC, Martin CA, Kim HS, Lai C, et al. Exosomal microRNA Differential Expression in Plasma of Young Adults with Chronic Mild Traumatic Brain Injury and Healthy Control. *Biomedicines* [Internet]. *Biomedicines*; 2021 [cited 2023 Jan 24];10:1–11. Available from: <https://pubmed.ncbi.nlm.nih.gov/35052715/>.

Waddington CH. The epigenotype. 1942. *Int J Epidemiol* [Internet]. *Int J Epidemiol*; 2012 [cited 2022 Oct 7];41:10–3. Available from: <https://pubmed.ncbi.nlm.nih.gov/22186258/>.

Willer, B., Leddy, J.J., 2006. Management of concussion and post-concussion syndrome. *Curr. Treat Options Neurol*.

Wyczechowska, D., Harch, P.G., Mullenix, S., Fannin, E.S., Chiappinelli, B.B., Jeansonne, D., et al., 2023. Serum microRNAs associated with concussion in football players. *Front Neurol* [internet]. *Frontiers Media SA 14* [cited 2024 Jan 11] Available from: </pmc/articles/PMC10151480/>.